Original Research Article

DOI: http://dx.doi.org/10.18203/2349-2902.isj20171150

Knowledge and attitude of dentists on bisphosphonates use in the UAE: a descriptive cross-sectional study

Kamis Gaballah^{1*}, Mawada Hassan²

¹Department of Oral and Maxillofacial Surgery, Ajman University, Ajman, United Arab Emirates

Received: 05 February 2017 **Accepted:** 02 March 2017

*Correspondence:

Dr. Kamis Gaballah,

E-mail: k.gaballah@ajman.ac.ae

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: The aim of this study was to assess the level of awareness and attitude of dentists practicing in the UAE about bisphosphonate intake and its dental implications.

Methods: A descriptive cross-sectional study using a 15 close-ended and fixed choice questions was performed on a representative sample of 502 dentists practicing in the UAE.

Results: The majority of the practitioners did not report treating patients on a bisphosphonate. Most of the participants were not familiar with the route of administration of the drug or the spectrum of side effects except for BRONJ which was regarded as the most common adverse effect. Furthermore, only a minority were aware of the standard treatment for patients on Bps.

Conclusions: The responses received not only highlight the lack of knowledge among practicing dentists regarding dental management of such cases but also urges the need for dissemination of guidelines for best practice in this field.

Keywords: Cross-sectional studies, Dentists, Disphosphonate, Drug-related side effects and adverse reactions

INTRODUCTION

Bisphosphonates (Bps) were first synthesized in 1868 in the industry. In the late 1960's Bps were shown to prevent the dissolution of hydroxyapatite of bone, the principal mineral component of bone, which made them a promising tool in the fight against bone metabolism disorders.¹ By the 1990's their actual mechanism of action was demonstrated with the launch of Fosamax® (Alendronate) by Merk.¹ On 2003, there were 17 million prescriptions for an alendronate worldwide making it among the top twenty most commonly prescribed drug worldwide.^{2,3} The popularity of the drug has risen over the last decade to become the treatment of choice for a number of serious disorders including; osteoporosis, cancer metastases to bone, hypercalcemia of malignancy, and multiple myeloma. The Bps inhibit osteoclastic bone

resorption through attachment to hydroxyapatite binding sites on bony surfaces, especially surfaces undergoing active resorption. The drug also reduces osteoclasts by decreasing their progenitor cell development and recruitment and also by promoting osteoclast apoptosis.²

However, the side-effect of Bps-related osteo-necrosis of the Jaw, BRONJ has emerged as an alarming consequence of drug use; between 2001-2007 with more than 2,400 confirmed cases reported in the USA alone.^{2,4} The BRONJ may represent the second epidemic of the phossy jaw which was reported between 1858 and 1906 with the similar clinical course of the current disease.⁸

To accommodate the growing number of osteonecrosis cases involving the maxilla and mandible associated with other antiresorptive (denosumab) and antiangiogenic

²Department of Clinical Dentistry, Ajman University, Ajman, United Arab Emirates

therapies, in addition to Bps, the Special Committee of American association of oral and maxillofacial surgeons has recommended that the nomenclature of BRONJ of the jaw be replaced with the term medication-related osteonecrosis of the jaw (MRONJ).^{6,7}

The dental and oral surgical intervention on the patient on various types of Bps therapy are susceptible to a higher rate of post-operative complications as compared to those not taking the medicine.^{3,4} Furthermore, patients on Bps therapy may seek dental care because of dental or jaw pain without any obvious signs of the complications of the therapy. If their symptoms do not resolve with routine dental and periodontal treatment, BRONJ must be considered as a differential diagnosis, even in the absence of exposed bone.⁵

The role of general dental practitioners (GDP) in prevention and early detection of BRONJ cases cannot be overemphasized. This study sought to assess the level of awareness and perceptions of practicing dentists in relation to bisphophonates and their associated risks, and how their knowledge influences the planning the management of patient taking such medications.

METHODS

This study was a descriptive cross sectional and involved a questionnaire which initially validated through multiple phases including experts review of the questions followed by a pilot study where 70 participants from Ajman, Sharjah and Dubai answered the questionnaire. The responses were revised for consistency of answers the negatively phrased questions. Finally, the principal component analysis was carried out to remove questions that measure the same variable. The revised questionnaire composed of 22 closed ended with fixed-choice questions. The proposal of the study was revised and approved by the research ethical committee at Ajman University. Then the questionnaire was distributed to a sample of 502 practicing dental practitioners. The questionnaire formed of the following components; 1) The frequency of patients on Bps seen by dental practitioners at regular basis, 2) The perception of practitioners of serious side effects of the drug, 3) The recognition of dental interventions that might be influenced by Bps therapy and 4) The knowledge of the optimal dental management of patients seek dental treatment during or after drug therapy.

The sample size was determined using the ministry of health records of the dentists currently practicing in the UAE. The number of target participants was calculated at 95% confidence interval with error margin less than 4%. The representative sample size was 499 dentists (Table 1). A random selection of dentists was invited to take part in the survey. The inclusion criteria were practice in the UAE and willingness to participate with no specific exclusion criteria.

Table 1: Sample stratification of the participants.

Stratification procedure									
Emirates	Stratification (ideal sample size)	No. of participants							
Abu- Dhabi	$(\frac{950}{2921}) \times 500 = 162$	160							
Dubai	$(\frac{1137}{2921}) \times 500 = 194$	190							
Sharjah	$(\frac{522}{2921}) \times 500 = 89$	83							
Ajman	$(\frac{124}{2921}) \times 500 = 21$	21							
Umm Al- quwain	$(\frac{15}{2921}) \times 500 = 5$	10							
RAK	$(\frac{108}{2921}) \times 500 = 18$	25							
Al- Fujairah	$(\frac{65}{2921}) \times 500 = 11$	13							
Total	500	502							

The data collected were entered and analyzed using software IBM SPSS Statistics version 20. Descriptive statistics were produced to classify the participants in relation to their specialties, working places and years of experience. The responses were compared in relationship to the participant data Chi square test. The statistical significance (p-value) was set at below 0.05 at 95 confidence rate.

RESULTS

The participants were divided according to their working place as following: 248 (49.4%) practice in private clinics, 105 (20.9%) in private hospitals, followed by 92 (18.3%) employed by public hospitals and 57 (11.4%) by public dental centers. More than two-thirds of the respondents 343 (n=343) had more than 5 years of experiences. The majority of the participants were general dental practitioners (302), the remaining were a quiet homogenous mix of the main dental specialties.

Frequency of patients on Bps receiving dental treatment

Two thirds of all participants didn't report treating any patients on Bps therapy followed by 19.9% (n=100) they had seen at least one patient on the medicine over the last year. Only 10 (1.99%) of those who claimed treated patients on Bps therapy were able to recall the brands of the drugs like Zometa, Fosamax, and Risedronate.

Awareness of the route of administration

A good number of participants (n=67, 13.3%) were not aware of the route of administration of the drug. The orthodontists gave the highest rate of unsatisfactory answers throughout the study. On the other hand, oral surgeons scored very high as compared to other dental disciplines (Table 2).

Indications of Bp therapy

While the majority of participants in the study (68.4%) identified, the osteoporosis being the commonest indication for Bps treatment, only 5.9% reported that cancer as another major indication to take the medicine

and only 1.6% mentioned other conditions like Paget's disease, multiple myeloma, fibrous dysplasia and osteogenesis imperfecta. The remaining a 15.7% of participants couldn't recall any indications for Bps prescription (Table 3).

Table 2: Knowledge on route of administration of the Bps.

Route of administration of Bps therapy										
			Administration	on						
			One satisfactory answer	>One satisfactory answer	Not satisfactory	Not aware	Total			
		Count	146	20	42	94	302			
	GDP	% within administration	60.1%	58.8%	62.7%	59.5%	60.2%			
		% of Total	29.1%	4.0%	8.4%	18.7%	60.2%			
		Count	20	8	3	9	40			
	Oral surgeon	% within administration	8.2%	23.5%	4.5%	5.7%	8.0%			
	J	% of Total	4.0%	1.6%	0.6%	1.8%	8.0%			
	Endodontists	Count	21	1	6	13	41			
		% within administration	8.6%	2.9%	9.0%	8.2%	8.2%			
C: -14		% of Total	4.2%	0.2%	1.2%	2.6%	8.2%			
Specialty	Prosthodontists	Count	22	3	5	11	41			
		% within administration	9.1%	8.8%	7.5%	7.0%	8.2%			
		% of Total	4.4%	0.6%	1.0%	2.2%	8.2%			
	Orthodontists	Count	25	0	7	14	46			
		% within administration	10.3%	0.0%	10.4%	8.9%	9.2%			
		% of Total	5.0%	0.0%	1.4%	2.8%	9.2%			
		Count	9	2	4	17	32			
	Others	% within administration	3.7%	5.9%	6.0%	10.8%	6.4%			
		% of Total	1.8%	0.4%	0.8%	3.4%	6.4%			
		Count	243	34	67	158	502			
Total		% within administration	100.0%	100.0%	100.0%	100.0%	100.0%			
		% of total	48.4%	6.8%	13.3%	31.5%	100.0%			

Table 3: The indication and potential side effects of the Bps treatment.

Indications of Bps and side effects of Bps therapy									
			Indications	Indications reported			Side effects of Bps		
			One	>One	Not	Satisfactory	Unsatisfactory	Not	Total
				indications	aware	answer	answer	aware	
	GDP		236	23	43	206	38	58	302
	ODI	% of Total	47.0%	4.6%	8.6%	41.0%	7.6%	11.6%	60.2%
	Oral surgeon	Count	26	11	3	28	3	9	40
	Oral surgeon	% of Total	5.2%	2.2%	0.6%	5.6%	0.6%	1.8%	8.0%
	Endodontists	Count	30	6	5	30	5	6	41
Specialty		% of Total	6.0%	1.2%	1.0%	6.0%	1.0%	1.2%	8.2%
Specialty	Prosthodontists	Count	30	4	7	23	9	9	41
		% of Total	6.0%	0.8%	1.4%	4.6%	1.8%	1.8%	8.2%
	Orthodontists	Count	33	2	11	17	14	15	46
		% of Total	6.6%	0.4%	2.2%	3.4%	2.8%	3.0%	9.2%
	Others	Count	19	3	10	22	0	10	32
	Oulers	% of Total	3.8%	0.6%	2.0%	4.4%	0.0%	2.0%	6.4%
Total		Count	374	49	79	326	69	107	502
		% within indications	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
		% of Total	74.5%	9.8%	15.7%	64.9%	13.7%	21.3%	100.0%

Common side effects of Bps therapy

About sixty percent of participants (n=314) regarded the BRONJ as the most serious side effect of the drug with a minority (n=12, 2.4%) added the oral ulcerations.

Approximately one-fifth of dentists (n=107) were not familiar with any side effects of Bps at all. The least awareness level reported by orthodontists (37%) (Table 3)

Table 4: The dental treatments that may be influenced by the Bps intake.

		Dental Rx that may be affected by BP-therapy							
			One relevant answer	More than one relevant answer	Not relevant answer	Not aware	Not asked	Not satisfactory	Total
	GDP	Count	113	93	12	30	40	14	302
_	ODF	% of Total	22.5%	18.5%	2.4%	6.0%	8.0%	2.8%	60.2%
	Oral surgeon	Count	6	23	1	1	6	3	40
_	Orar surgeon	% of Total	1.2%	4.6%	0.2%	0.2%	1.2%	0.6%	8.0%
	Endodontists	Count	16	14	1	4	2	4	41
Cnacialty -		% of Total	3.2%	2.8%	0.2%	0.8%	0.4%	0.8%	8.2%
Specialty -	Prosthodontists	Count	11	13	5	3	9	0	41
_		% of Total	2.2%	2.6%	1.0%	0.6%	1.8%	0.0%	8.2%
	Orthodontists	Count	19	11	4	5	6	1	46
_	Orthodontists	% of Total	3.8%	2.2%	0.8%	1.0%	1.2%	0.2%	9.2%
	Others	Count	4	13	1	6	7	1	32
Others		% of Total	0.8%	2.6%	0.2%	1.2%	1.4%	0.2%	6.4%
		Count	169	167	24	49	70	23	502
Total		% of Total	33.7%	33.3%	4.8%	9.8%	13.9%	4.6%	100.0 %

Table 5: The dental treatment of patients on Bps therapy.

Dental treatment of	of patients on Bps	therapy				
			Dental treatments			
			Standard practice	Substandard practice	Not aware	Total
		Count	11	202	89	302
	GDP	% within treatments	64.7%	61.2%	57.4%	60.2%
		% of total	2.2%	40.2%	17.7%	60.2%
		count	2	30	8	40
	Oral surgeon	% within treatments	11.8%	9.1%	5.2%	8.0%
		% of total	0.4%	6.0%	1.6%	8.0%
	Endodontists	count	1	30	10	41
		% within treatments	5.9%	9.1%	6.5%	8.2%
C:-14		% of total	0.2%	6.0%	2.0%	8.2%
Specialty	Prosthodontists	count	2	27	12	41
		% within treatments	11.8%	8.2%	7.7%	8.2%
		% of total	0.4%	5.4%	2.4%	8.2%
	Orthodontists	count	1	25	20	46
		% within treatments	5.9%	7.6%	12.9%	9.2%
		% of total	0.2%	5.0%	4.0%	9.2%
		count	0	16	16	32
	Others	% within treatments	0.0%	4.8%	10.3%	6.4%
		% of total	0.0%	3.2%	3.2%	6.4%
Total		count	17	330	155	502
		% of total	3.4%	65.7%	30.9%	100.0%

Table 6: Dentist view on discontinuation of the Bps.

Discontinuation of Bps therapy										
				Discontinuation			If Yes			
				Yes	I don't Know	Total	5-10 days	1 month	3 months	Total
	count count		77	78	147	302	21	19	38	78
	GDP	% of total	15.3%	15.5%	29.3%	60.2%	14.5%	13.1%	26.2%	53.8%
	Oral surgaan	count	16	17	7	40	0	1	16	17
	Oral surgeon	% of total	3.2%	3.4%	1.4%	8.0%	0.0%	0.7%	11.0%	11.7%
	Endodontists	count	5	18	18	41	4	4	10	18
Cmanialty		% of total	1.0%	3.6%	3.6%	8.2%	2.8%	2.8%	6.9%	12.4%
Specialty	Prosthodontists	count	7	15	19	41	8	2	5	15
		% of total	1.4%	3.0%	3.8%	8.2%	5.5%	1.4%	3.4%	10.3%
	Outhodontists	count	7	10	29	46	4	2	4	10
	Orthodontists	% of total	1.4%	2.0%	5.8%	9.2%	2.8%	1.4%	2.8%	6.9%
	Others	count	5	7	20	32	3	2	2	7
	Ouleis	% of total	1.0%	1.4%	4.0%	6.4%	2.1%	1.4%	1.4%	4.8%
Total	TD-4-1		117	145	240	502	40	30	75	145
Total		% of total	23.3%	28.9%	47.8%	100.0%	27.6%	20.7%	51.7%	100.0%

Dental interventions affected by Bp therapy

When asked about the impact of Bp therapy on dental treatment, the responses obtained from various participant were not consistent with one quarter (n=124) believe the only extraction of teeth is adversely affected by the Bps intake. However, 15.9% of the participant expanded their responses to include periodontal treatment and insertion of implants and others (11.9) added the orthodontic treatment to the previous list. On the other hand, around one third of participants (n=166) were not aware of any relationship between the Bps therapy and dental treatment interventions.

Dental management of patients on Bps therapy

Thirty percent of respondents were not aware of the 'standard' dental treatments for a patient on Bps treatment. The majority (65.7%) of participants reported substandard treatment plans for such a patient, these include: planned surgical tooth extraction versus closed method (35.1%), routine extraction of teeth without any additional measures (10.8%), prescribing antibiotics and retain the tooth in situ (9.6%), or performing conventional endodontic treatment to affected tooth (6.6%). The answers of only 17 (3.4%) divulge a standard treatment for patients required dental extraction during their treatment with Bps (Table 5). The responses of nearly half of the participating dentists (47.8%) showed a lack of awareness of issues of discontinuation of Bps prior to dental and surgical intervention. However, about one third of dentists suggested to stop the drug for; 5 days (4%), 10days (4%), 1 month (4), or 3 months (14.9%) prior to the commencement of dental treatment. Around one quarter of participants (23.3%) do not consider stopping the drug pre-operatively. When asked about using alternative medications instead of Bps or altering the dose of such therapy, almost half of the participants (48.2%) were ready to adopt such an approach (Table 6).

International guidelines on dental management of patient on Bps therapy

Nearly two thirds of participants irrespective of their specialties (n=337) were not aware of any specific guidelines for dental management of patients taking Bps. Around one third (n=175) of participants would prescribe antibiotics to patients on Bps regardless the dental interventions as compared to 3.8% of who opposed antibiotics prescription in all the dental interventions. On the other hand, 40.6% who recommended the antibiotics when only surgical extraction is planned for such patients. Finally, most of the participants (89.2%, n=448) were keen to attend updated CME courses on the dental management of patients on Bps.

DISCUSSION

The Bps group of medications have proven effective in treating many conditions. They were introduced to the middle-east and Gulf region quiet recently. However, their prescription showed a significant spread due to the demand of local patients and the relative absence of tight authoritive restriction. The current study obviously has the limitations of the questionnaire-based survey and may not precisely reflect the real practice of the participants. The authors consider the possibility of response distortion as participants attempt to create a positive impression that is aware of all aspects of the issue. The tendency to respond positively and not revealing the weakness can adversely influence the outcome of this survey. However, the study of current practice usually starts with such surveys to understand the awareness and attitude of the dentists till it can be further verified by investigation of patient records when a sufficient number of the later are available for review and study.

Despite the noticeable expansion of the drug prescription in the region, about two-thirds of participants did not report treating any patient using the medicine. This may, in part, indicate that Bps intake is not yet among the routine case history checklist of dental practice in this region. Furthermore, this reflects the lack of awareness of the drug's effects and side-effects, more importantly, the clinical indications for their use. This notion is supported by the failure of a good proportion of respondents to recall any indication for Bps use with nearly three quarters only identified the osteoporosis as a sole indication. However, this positive response from the majority of participants was not matched with their failure to report treating any patient using the drug considering the relatively high incidence of osteoporosis cases in this region and the increasing trend in prescribing Bps to treat the condition. Failure of respondents to report any additional indications for the drug use implies that dentist may not to anticipate that their dental patients might be using this medicine for various problems thus will not enquire about the intake of Bps in such cases. More than half of the dentists reported that they are aware of BRONJ of side effect of the drug which can be compared favourably to the number of healthcare professionals aware of BRONJ as a side-effect in North Wales.9

This study also showed that two-thirds of the practicing dentists are aware of the dental interventions that might be troubled by Bps therapy such as; oral surgery, periodontal therapy, and orthodontics. The awareness rate reported here was higher than that reported by dentists in South Korea (56.5%) (10, 11), in Canada (60%) (12-16) and physicians in Saudi Arabia (31%) (17,18). Dentoalveolar surgery may cause or exacerbate an existing BRONJ in patients during or after Bps therapy. Although this is more prevalent in those taking intravenous Bps, it has also been reported in those taking these drugs orally over long periods of time. Concomitant risk factors include poorly controlled diabetes mellitus and other immunocompromised states, concurrent use of corticosteroids, chemotherapeutic drugs, radiotherapy, advanced age, alcohol abuse and smoking.

It is estimated that the majority of BRONJ cases are preceded by dental surgical treatment, with twice as many occurring in mandible as in maxilla. It was suggested that Bp treatment should be postponed during active dental treatment is concluded to minimize the risk of BRONJ development. ^{2,3,9} Local risk factors include the presence of mandibular tori, periodontal disease, periapical disease and recent trauma. ^{11,19,20}

In a South Korean study, involving 226 dentists, around half of the respondents were aware of BRONJ, with the practitioners with fewer than 5 years' experience were reported to be significantly more aware of the condition

than those qualified for more than 5 years. 11 This reflects that younger dentists know more about the medicine as compared to senior counterparts and this was also supported by the current study. Unfortunately, only minority of participants were familiar with the standard treatment of patients on Bps require tooth extraction procedure. Moreover, the highest level of substandard practice was mainly reported by oral surgeons who are expected to manage the oral complications seen on the patient on Bps. A referral of patients for comprehensive oral evaluation before commencing Bps treatment should be considered by the prescribing physicians.²¹ Once patient started the Bps treatment, a great emphasis on oral hygiene measured should be exercised to minimise risk factors for infection. Regular prophylactic chlorhexidine rinse and modified treatments such as coronal amputation and endodontic treatment of retained roots are advised for non-restorable teeth. If surgery is unavoidable, focus on minimally invasive procedures with proper sterile technique and effective empirical antibiotic therapy.²²⁻²⁴

This study also raises the issue of drug holiday preoperatively, most of the respondents did not show any updated or evidence based knowledge on this option. Many researchers questioned the value of stopping Bps pre-operatively as the drug tend to remain bound to the bone for years after the administration. On the contrary, others suggested considering interrupting Bps treatment for three months prior to surgery and restarting after bone healing. This suggestion has not been supported by the ADA. According to the latter, no evidence to date has confirmed that drug holidays are effective in preventing BRONJ without increasing the skeletally-related risks of low bone mass during treatment. Late of the service of the stellar of the skeletally-related risks of low bone mass during treatment.

The current study analysis reveals that most of the participants were not familiar with any dental guidelines for treating patients with Bps therapy that highlight the importance of CME courses and establishing local authoritive guidance in this area.

CONCLUSION

An increasing number of dental patients are currently undergoing a form of Bps therapy and according to this descriptive survey, the dentists do not seem to be aware of this. It is also apparent that those dentists who are aware of their patients' drug regimens are not always aware of all potential side effects of these medications and the impact of poorly planned dental treatment. The responses received not only highlight the lack of knowledge among dentists regarding Bps and their dental implications but also reinforce the value of dissemination of guidelines for best practice in this field.

Funding: No funding sources Conflict of interest: None declared

Ethical approval: The study was approved by the

institutional ethics committee

REFERENCES

- 1. Fleisch H. Development of bisphosphonates. Breast Cancer Res. 2001;4(1):30-4.
- 2. Krueger CD, West PM, Sargent M, Lodolce AE, Pickard SA. Bisphosphonate-induced osteoporosis of the jaw. Ann Pharmacother. 2007;41(1):276-84.
- 3. British Dental Association, 2008. Bisphosphonates. Fact File. London: BDA.
- 4. Arrain Y, Masud T. Recent recommendations on bisphosphonate-associated osteochemonecrosis of the jaws. Dent Update. 2008;35(1):238-42.
- 5. Mosaferi H, Fazlyab M, Sharifi S, Rahimian S. Bisphosphonate-induced osteonecrosis of the maxilla resembling an infectious endodontic lesion. Iran Endod J. 2016;11(1):67-70.
- Ruggiero SL, Dodson TB, Assael LA. American Association of Oral and Maxillofacial Surgeons position paper on bisphosphonate-related osteonecrosis of the jaws- 2014 update. J Oral Maxillofac Surg. 2014;72(6):1-26.
- 7. Yamashita J, McCauley LK. Antiresorptives and osteonecrosis of the jaw. J Evid Based Dent Pract. 2014;12(233):233-47.
- 8. Masson DR, O'Calloghan E, Seager M. The knowledge and attitudes of North Wales health care professionals to bisphosphonate associated osteochemonecrosis of the jaws. J Disability Oral Health. 2009;10(4):175-83.
- Alhussain A, Peel S, Dempster L, Clokie C, Azarpazhooh A. Knowledge, practices, and opinions of Ontario dentists when treating patients receiving bisphosphonates. J Oral Maxillofac Surg. 2015;73(6):1095-105.
- Al-Mohaya MA, Al-Khashan HI, Mishriky AM, Al-Otaibi LM. Physicians' awareness of bisphosphonates-related osteonecrosis of the jaw. Saudi Med J. 2011;32(8):830-35.
- 11. Maha A. Al-Mohaya. Hesham I. Physicians' awareness of bisphosphonates-related osteonecrosis of the jaw. Saudi Med. 2011;32(8):830-35.
- Sirmahan C, Firat S, Cengizhan K. Bisphosphonates and Bone, Orthopedic Surgery. 1st ed. Intechopen. 2012.
- 13. Russell RG. Bisphosphonates: The first 40 years. Bone. 2011;49(1):2-19.

- 14. Giger EV. Castagner B and Leroux JC. Biomedical applications of bisphosphonates. J Control Release. 2013;167(1):175-88.
- 15. Xu XL, Gou WL, Wang AY, Wang Y, Guo QY, Lu Q, et al. Basic research and clinical applications of bisphosphonates in bone disease: what have we learned over the last 40 years? J Trans Med. 2011;11(303):1-8.
- 16. Lim V, Clarke BL. New therapeutic targets for osteoporosis: beyond denosumab. Maturitas. 2012;73(1):269-72.
- Basso FG. Turrioni AP. Hebling J, de Souza CC. Effects of zoledronic acid on odontoblast-like cells. Arch Oral Biol. 2013;58(1):467-73.
- 18. Abela S, Chotai M, Bister D. What you need to know about bisphosphonates: an overview and general recommendations for orthodontic treatment. J Orthod. 2012;39(19):186-92.
- Saldanha S, Shenoy VK, Eachampati P, Uppal N. Dental implications of bisphophonate-related osteonecrosis. Gerodontology Society and John Wiley and Sons A/S. Gerodontology. 2012;29(3):177-87.
- 20. Lewis DM. Clinical: Bisphosphonates, phossy jaw, and BRONJ. ODA J. 2011;102(5):36-9.
- Tsao C, Darby I, Ebeling PR. Oral health risk factors for bisphosphonate-associated jaw osteonecrosis. J Oral Maxillofac Surg. 2013;71(8):1360-6.
- 22. Lee JK, Kim KW, Choi JY, Moon SY, Kim SG, Kim CH, et al. Bisphosphonates-related osteonecrosis of the jaw in Korea: a preliminary report. J Korean Assoc Oral Maxillofac Surg. 2013;39(1):9-13.
- 23. Diab DL, Watts NB. Bisphosphonate drug holiday: who, when and how long. Ther Adv Musculoskel Dis. 2013;5(3):107-11.
- 24. Compston J, Bilezikian J. Bisphosphonate therapy for osteoporosis: The long and short of it. J Bone Miner Res. 2013;27(2):240-2.

Cite this article as: Gaballah K, Hassan M.

Knowledge and attitude of dentists on bisphosphonates use in the UAE: a descriptive cross-sectional study. Int Surg J 2017;4:1398-404.