pISSN 2349-3305 | eISSN 2349-2902

## **Review Article**

DOI: https://dx.doi.org/10.18203/2349-2902.isj20253737

# Latissimus dorsi flap for breast reconstruction

## Juan Carlos García Zamora\*

Department of General and Bariatric Surgery, Private Practice, Guadalajara, Jalisco, Mexico

Received: 16 October 2025 Accepted: 31 October 2025

# \*Correspondence:

Dr. Juan Carlos García Zamora, E-mail: jucagaza@hotmail.com

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### **ABSTRACT**

The latissimus dorsi flap (LDF) is a reliable and time-tested autologous option for breast reconstruction following mastectomy or breast-conserving surgery. It can be harvested as a myocutaneous or muscle-only flap, with or without fat grafting or implant placement, allowing tailored reconstruction based on patient anatomy, comorbidities, and aesthetic goals. The LDF is especially advantageous in patients who are not suitable candidates for abdominal-based free flaps or who present with compromised local tissues after radiation or failed reconstruction. Recent refinements, such as immediate fat augmentation and muscle-sparing techniques, have expanded its reconstructive versatility and improved donor-site outcomes. The most common complications are donor-site related, with seroma formation being the predominant issue, reported in up to 40% of cases. Other complications include hematoma, wound dehiscence, infection, and partial flap necrosis, though total flap loss is rare. Functional morbidity, particularly mild shoulder weakness or limited range of motion, may occur but generally has minimal impact on daily activities. Techniques preserving the thoracodorsal nerve and minimizing muscle harvest significantly reduce postoperative morbidity. Overall, the LDF provides consistent aesthetic and functional results, making it a valuable and versatile reconstructive option. Its safety, reliability, and adaptability-particularly when integrated with fat grafting-ensure its continued role in contemporary breast reconstruction.

**Keywords:** Latissimus dorsi flap, Breast reconstruction, Donor-site morbidity, Seroma, Shoulder function, Fat grafting, Muscle-sparing technique, Reconstructive surgery

## INTRODUCTION

The latissimus dorsi flap (LDF) is a well-established autologous option for breast reconstruction, utilized in both immediate and delayed settings following mastectomy or breast-conserving surgery. The flap can be harvested as either a myocutaneous or muscle-only flap, with or without adjunctive fat grafting or implant placement, depending on the reconstructive requirements and patient-specific factors.<sup>1-8</sup>

LDF is particularly valuable in patients who are not candidates for abdominal-based free flaps (e.g., due to insufficient abdominal tissue, prior abdominal surgery, or comorbidities precluding microsurgery) and in those with poor local tissue quality, such as post-radiation or failed implant-based reconstruction.<sup>1,3,5,8</sup> The technique is

reliable, with consistently low rates of major complications and flap loss, and is associated with high patient satisfaction and favorable aesthetic outcomes.<sup>2-8</sup>

The flap may be used for: Total breast reconstruction, either as a stand-alone autologous procedure or in combination with an implant or tissue expander to achieve adequate volume.<sup>2,3,5-8</sup> Partial breast reconstruction to address defects following breast-conserving therapy, especially in low-volume, nonptotic breasts.<sup>9</sup> Salvage reconstruction after failed implant or free flap procedures.<sup>3,8</sup>

Technical modifications, such as immediate fat grafting (fat-augmented LDF), have been developed to overcome the inherent volume limitations of the latissimus dorsi muscle and improve contour and softness, further

enhancing aesthetic results and patient satisfaction.<sup>1,7</sup> The choice between muscle-only and myocutaneous flaps may be guided by the need for skin supplementation and donor site considerations, with muscle-only flaps offering shorter operative times and improved donor site satisfaction when skin replacement is not required.<sup>4,7,8</sup>

Common complications include seroma formation at the donor site, wound dehiscence, and, less frequently, partial flap necrosis; however, serious adverse events and total flap loss are rare.<sup>5</sup> Functional outcomes are generally excellent, with minimal long-term impairment of arm or back function.<sup>6</sup>

### **COMPLICATIONS**

The LDF procedure is associated with several well-characterized complications, which can be categorized as donor-site, flap-related, and functional morbidities.

Donor-site complications are common and include seroma formation, which is consistently reported as the most frequent issue, with rates ranging from approximately 26% to over 40% depending on patient population and flap technique. Seromas often require aspiration or drainage, typically at the donor site. Hematoma is less common but still notable, with rates around 2-10% and a higher incidence compared to other flap types in head and neck reconstruction. Wound dehiscence and surgical site infection occur in 7-8% of cases, and partial skin or nipple necrosis is seen in about 7% of patients. Poor wound healing is also reported, with an incidence of 8.6% in head and neck reconstruction. Flap loss due to necrosis is rare, occurring in approximately 2% of cases.

Functional donor-site morbidity primarily involves shoulder dysfunction. Quantitative studies demonstrate a significant reduction in shoulder adduction and extension strength (17-21% loss at 12 months), although range of motion and ability to perform activities of daily living are generally preserved. Subjective discomfort at the donor site is reported by up to 41% of patients, with most experiencing minimal impact on daily activities but greater difficulty with sports or artistic activities. The degree of functional impairment is influenced by the extent of muscle harvested and whether the thoracodorsal nerve is preserved; muscle-sparing and nerve-preserving techniques are associated with lower morbidity. Sensory disturbances at the donor site are also common and may increase over time.

Other complications include hematoma/seroma at the donor site, which is more prevalent with LDF s compared to anterolateral thigh flaps.<sup>17</sup> The risk of complications such as seroma and prolonged drainage is higher with larger flap sizes and in breast reconstruction settings.<sup>2</sup> Patient factors such as obesity, smoking, and prior radiation do not appear to significantly increase donor-

site complication rates, although obese patients may have a higher risk of mastectomy skin flap necrosis. 18

In summary, the most frequent complications of the LDF procedure are donor-site seroma, hematoma, wound dehiscence, infection, partial necrosis, and functional shoulder impairment, with the severity and incidence influenced by surgical technique and patient factors.<sup>18</sup>

#### **CONCLUSION**

The LDF remains a dependable and versatile option for breast reconstruction, offering predictable outcomes, high flap survival, and broad applicability across a range of reconstructive scenarios. Its reliability, even in patients with prior radiation, comorbidities, or failed previous reconstructions, underscores its enduring relevance in modern reconstructive surgery. Technical refinements-such as fat-augmented and muscle-sparing variations—have further improved both aesthetic outcomes and donor-site morbidity, making the procedure more adaptable to individual patient needs.

While donor-site complications, particularly seroma formation, remain the most frequent issue, these are typically minor and manageable with appropriate drainage and postoperative care. Functional morbidity, primarily related to shoulder strength and range of motion, is generally mild and well tolerated, especially when nerve-preserving and limited-harvest techniques are employed. Serious complications such as flap loss or significant disability are uncommon.

In conclusion, the latissimus dorsi flap continues to serve as a cornerstone in autologous breast reconstruction. Its balance of safety, versatility, and aesthetic potential-particularly when combined with modern adjuncts like fat grafting-makes it an invaluable option for both primary and salvage reconstruction, with excellent long-term functional and cosmetic results.

Funding: No funding sources Conflict of interest: None declared Ethical approval: Not required

# **REFERENCES**

- 1. Abdou SA, Charipova K, Song DH. Modern approaches to pedicled latissimus dorsi flap breast reconstruction with immediate fat transfer. Clin Plast Surg. 2023;50(2):259-65.
- 2. Liu Q, Li W, Wu X, Liang X, Pinghua H, Yali C. Immediate breast reconstruction using latissimus dorsi muscular flap: A retrospective study of Chinese patients with breast cancer. Medicine. 2021;100(24):e26175.
- 3. Zheng S, Hao S, Chen J, Yingying Z, Benlong Y, Xiaoyan H, et al. Latissimus dorsi flap-the main force in breast reconstruction for breast tumor in Chinese population. Front Oncol. 2023;13:1159073.

- 4. Cogliandro A, De Bernardis R, Calabrese G, Rosa S, Valeria P, Barbara C, et al. Muscular versus myocutaneous latissimus dorsi flap in delayed breast reconstruction: A comparative study of postoperative outcomes. Aesthetic Plast Surg. 2025;NA.
- Banys-Paluchowski M, Brus L, Krawczyk N, Sophie VK, Maria LG, Nana B, et al. Latissimus dorsi flap for breast reconstruction: A large single-institution evaluation of surgical outcome and complications. Arch Gynecol Obstet. 2024;309(1):269-80.
- 6. Zheng H, Zhu G, Guan Q, Wei F, Xiang L, Mancheng Y, et al. A retrospective study of latissimus dorsi flap in immediate breast reconstruction. Front Oncol. 2021;11:598604.
- Tomita K, Taminato M, Kubo T. Total breast reconstruction with a fat-augmented latissimus dorsi flap: A comparative study between muscle and myocutaneous flaps. J Plast Reconstruct Aesthet Surg. 2023;83:250-7.
- Bucaria V, Giudice G, Boccuzzi A, Ilaria C, Michele M, Rossella E. "No-back-scar" latissimus dorsi muscle flap and tissue expander: A valuable strategy for secondary breast reconstructions. Aesthetic Plast Surg. 2024;48(11):2098-107.
- 9. Mericli AF, Szpalski C, Schaverien MV, Jesse CS, David MA, Patrick BG, et al. The latissimus dorsi myocutaneous flap is a safe and effective method of partial breast reconstruction in the setting of breast-conserving therapy. Plast Reconstruct Surg. 2019;143(5):927e-35.
- 10. Banys-Paluchowski M, Brus L, Krawczyk N, Sophie VK, Maria LG, Nana B, et al. Latissimus dorsi flap for breast reconstruction: A large single-institution evaluation of surgical outcome and complications. Arch Gynecol Obstet. 2024;309(1):269-80.
- Arikawa M, Miyamoto S, Fujiki M, Takuya H, Azusa O, Minoru S. Comparison of donor site drainage duration and seroma rate between latissimus dorsi musculocutaneous flaps and thoracodorsal artery perforator flaps. Ann Plast Surg. 2017;79(2):183-5.

- 12. Wang Y, Liang C, Lin Q, Lin F, Libo L, Haoran J, et al. Donor-site morbidity and aesthetic outcomes in patients undergoing head and neck reconstruction with anterolateral thigh or latissimus dorsi flaps: A systematic review and meta-analysis. J Cranio-Maxillo-Facial Surg. 2025;53(10):1892-903.
- 13. Højvig JH, Henriksen M, Bartholdy CR, Bonde CT. Donor-site morbidity following breast reconstruction with a latissimus dorsi flap: A prospective study. J Plast Reconstruct Aesthet Surg. 2022;75(7):2205-10.
- 14. Lee KT, Mun GH. A systematic review of functional donor-site morbidity after latissimus dorsi muscle transfer. Plast Reconstruct Surg. 2014;134(2):303-14.
- 15. Oberhofer HM, Samant SS, Swan CC, Erin MW, Ellen SS, Mark ML, et al. Objective comparison of donor-site morbidity following full and thoracodorsal nerve-preserving split latissimus dorsi flaps. Plast Reconstruct Surg. 2022;149(5):966e-71e.
- 16. Gatto A, Parisi P, Brambilla L, Ilaria S, Annarita V, Federico LT, et al. Thoracodorsal artery perforator flap, muscle-sparing latissimus dorsi, and descending branch latissimus dorsi: A multicenter retrospective study on early complications and meta-analysis of the literature. J Plast Reconst Aesthet Surg. 2022;75(11):3979-96.
- 17. Yezhelyev M, Duggal CS, Carlson GW, Losken A. Complications of latissimus dorsi flap breast reconstruction in overweight and obese patients. Ann Plast Surg. 2013;70(5):557-62.
- 18. Steffenssen MCW, Kristiansen AH, Damsgaard TE. A systematic review and meta-analysis of functional shoulder impairment after latissimus dorsi breast reconstruction. Ann Plast Surg. 2019;82(1):116-27.

**Cite this article as:** Zamora JCG. Latissimus dorsi flap for breast reconstruction. Int Surg J 2025;12:xxx-xx.