Original Research Article

Epidemiological profile of cancer patients in surgery ward of a tertiary teaching hospital in the Vindhya region with special reference to high proportion of carcinoma gall bladder

Dhanwantari Shukla*, Vishnu Kumar Patel, Akash Singh Chhari, Chandra Shekhar Dubey, Rishi Kumar Garg, Singh A. P., Bais P. S., Gaharwar A. P. S.

Department of Surgery, Shyam Shah Medical College, Rewa, Madhya Pradesh, India

Received: 25 April 2016

Accepted: 03 June 2016

*Correspondence:

Dr. Dhanwantari Shukla,

E-mail: 6487sd@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Cancer is on rise. WHO estimates that by 2030, the incidence of cancer will increase by 68% as compared to 2012. This study evaluates epidemiological profile of Cancer in the Vindhya region. Design of this study was combination of retrospective and prospective study.

Methods: All cancer patients admitted in surgery ward of S. S. medical college, Rewa between 2010 to 2014 were included in the study. Data regarding age, gender, religion, socio-economic status, residence, dietary habits, addiction and site of cancer as well as mortality pattern of cancer was tabulate, analyzed and compared.

Results: 1186 cases were included in this study which constituted 3.02% of total admissions in surgical ward. 50.25% were males, 49.75% were females. Most common age groups were middle age group i.e. 35-64 years (25.30% males, 32.29% females, total 57.59%) and geriatric (19.48% males, 11.38% females, total 30.86%). 95.78% were Hindus, 4.22% were Muslims. 70.40% were BPL, 29.60% were APL. 67.20% belonged to rural areas, 32.80% were from urban areas. 75.89% were vegetarians, 24.11% were non-vegetarians. 25.97% were addicted to tobacco alone, 1.52% to alcohol alone and 18.63% to both tobacco and alcohol. Breast cancer was the most prevalent cancer (19.06%), followed by cancer gall bladder (10.55%). Among males, colorectal cancer was the commonest (11.24%), gall bladder was the 4th commonest (8.22%). Among females, breast was the commonest (37.12%) and gall bladder was the 2nd commonest (12.88%). Gall bladder cancer was the commonest (13.93%) in geriatric age group, 2nd commonest (9.37%) in the middle age group and 5th commonest in the young age group (7.63%). 18.38% of all and 46.40% of gall bladder cancer presented with metastasis. 91 cancer deaths were recorded accounting for 3.64% of all deaths and the case fatality rate of cancer was 7.67%. Gall bladder cancer was the commonest cause of Cancer deaths overall (14.29%) and in females (20.00%) and 2nd commonest among males (10.70%) after Ca stomach (12.50%). Cancer gall bladder was the commonest cause of death in middle age (16.34%) and 2nd commonest in geriatric age group (15.63%).

Conclusions: Cancer affects all genders, ages and demographic profiles. Cancer of gall bladder has emerged as a major cause of morbidity and mortality in this study.

Keywords: Cancer, Epidemiological, Gall bladder, Mortality

INTRODUCTION

Cancer is on rise. WHO estimates that by 2030, the incidence of cancer will increase by 68% as compared to

2012.¹ Neoplasm which can be defined as excessive and unregulated proliferation of cells independent of physiological growth-stimulatory stimuli is said to have become malignant when it attains a property of invading

and destroying adjacent structures and spreading to distant sites (metastasis) to cause death.² Cancer has a 5 millennia old history but it continues to haunt human societies and perplex clinicians with its wide variety of clinical course and often fatal outcome. Cancer is associated with a tremendous socio-economic as well as psychological burden. The annual incidence of Cancer is 14 million and it kills 8.2 million people annually.³ 1 in every 8 deaths is due to cancer.⁴ More than a million Indians are diagnosed with cancer each year while 6-7 lakhs of them lose their lives annually. The incidence and mortality both have increased by about 50% over past 2 decades.⁵ The greatest strides in the field of oncology have been made by epidemiological studies which helped in identifying the carcinogenic factors subsequently leading to discovery of the therapeutic modalities. 80-90% of all cancers are believed to arise from environmental and lifestyle risk factors.⁶ The world of Cancer management has seen a revolutionary change over the past century. From the days of clinical diagnosis, we have come to the era of tissue diagnosis and conventional imaging modalities like X-ray and USG are being supplemented with advanced imaging techniques like CT, MRI, PET and other radioisotope-labeled studies. Cancer therapeutics involves multiple disciplines like surgery, chemotherapy, radiotherapy and immunotherapy. Cancer of gall bladder is often asymptomatic or vaguely symptomatic, prompting patients to present at a stage where curative treatment is no more an option. Gall bladder cancer is the commonest biliary tract malignancy and is particularly common in North and Central India.¹³ This study describes the epidemiological profile of Cancer patients admitted in a tertiary teaching Hospital in the Vindhya region with regards to various demographic parameters and the organs involved and highlights the high prevalence and mortality of cancer of gall bladder.

The objective of this study was to know the overall incidence of malignancy and its distribution in relation to age, sex, occupation, religion, socio-economic status, residence, dietary habits, addiction and the stage of disease in which the diagnosis was made, to study the change in the trend of cancer incidence and mortality over a study period of 5 years. And study mortality in various malignancies, to highlight the high prevalence and mortality of cancer of gall bladder.

METHODS

This study was carried out on 1186 patients of cancer admitted in the surgical ward of S. S. medical college, Rewa between 1st January 2010 to 31st December 2014. Records of cancer patients admitted between 1st January 2010 - 31st May 2014 were obtained from Central registration of surgery department and patients diagnosed with cancer from 1st June 2014 to 31st December 2014 were examined individually. Age, gender, residence, occupation, addiction, socio-economic status, dietary habits, presenting complaints, history, examination, investigation, treatment and outcome were recorded meticulously in proforma/master chart for systematic tabulation, observation and statistical analysis.

RESULTS

It is evident from the Table 1 that incidence of cancer was almost equal in males (50.25%) and females (49.75%). Maximum (57.59%) of the patients were from the middle age group (35-64 years) in both genders. The proportion in geriatric age group (65+years) was 30.86% and was the 2nd commonest age group in both genders. The proportion of patients in young adults (15-34 years) was 11.05% and that of pediatric population (0-14 years) was 0.50%.

Table 1: Distribution of patients according to age and gender (n = 1186).

Age (years)	Male		Female		Total	
	Number	%	Number	%	Number	%
0-14	2	0.17	4	0.34	6	0.50
15-34	63	5.32	68	5.73	131	11.05
35-64	300	25.30	383	32.29	683	57.59
65+	231	19.48	135	11.38	366	30.86
Total	596	50.25	590	49.75	1186	100.00

It is evident from the Table 2 that 95.78% of the patients were Hindus, 4.22% were Muslims. About two-thirds (67.20%) belonged to rural areas, about one-third (32.80%) belonged to urban areas. The percentage of BPL (below poverty line) patients was 70.40%, rest

29.60% were APL (above poverty line). About three-fourths (75.89%) patients were vegetarians, one-fourths (24.11%) were non-vegetarians. 25.97% patients were addicted to tobacco alone, 1.52% to alcohol alone, 18.63% to both tobacco and alcohol, 53.88% were not addicted.

Table 2: Distribution of patients according to religion, residence, socio-economic status, dietary habits and addiction (n = 1186).

Characteristic		Number	Percentage
Religion	Hindus	1136	95.78
	Muslims	50	4.22
	Total	1186	100.00
Residence	Urban	389	32.80
	Rural	797	67.20
	Total	1186	100.00
Socio-economic status	APL	351	29.60
	BPL	835	70.40
	Total	1186	100.00
Dietary habits	Vegetarians	900	75.89
	Non-vegetarians	286	24.11
	Total	1186	100.00
Addiction	Tobacco alone	308	25.97
	Alcohol alone	18	1.52
	Tobacco+alcohol	221	18.63
	No addiction	639	53.88
	Total	1186	100.00

Table 3: Gender wise distribution of 10 commonest cancers (n = 1186).

Males			Females		
Site	No.	Percentage	Site	No.	Percentage
Colorectal	67	11.24	Breast	219	37.12
Prostate	61	10.24	Gall bladder	76	12.88
Penis	61	10.24	Colorectal	43	7.28
Gall Bladder	49	8.22	Cervix	39	6.61
Stomach	45	7.55	Stomach	33	5.59
Skin+soft tissue	42	7.05	Cholangiocarcinoma	24	4.06
Periampullary	41	6.88	Skin+soft tissue	22	3.73
Oral cavity	41	6.88	Periampullary	19	3.22
Hepatocellular	27	4.53	Hepatocellular	19	3.22
Urinary bladder	24	4.03	Ovary	19	3.22
10 commonest	458	76.84	10 commonest	513	86.94
Others	138	23.16	Others	77	13.06
Total	596	100.00	Total	590	100.00

It is evident from the table that among males, the 10 commonest cancers were those of colorectum (11.24%), prostate (10.24%), penis (10.24%), gall bladder (8.22%), stomach (7.55%), skin+soft tissue (7.05%), periampullary (6.88%), oral cavity (6.88%), hepatocellular (4.53%) and urinary bladder (4.03%). Among females, the 10 commonest cancers were those of breast (37.12%), gall bladder (12.88%), colorectum (7.28%), cervix (6.61%), stomach (5.59%), cholangio (4.06%), skin+soft tissue (3.73%), periampullary (3.22%), hepatocellular (3.22%) and ovary (3.22%).

It is evident from the Table 4 that in the pediatric age group, Wilm's tumour of kidney was the commonest cancer accounting for 66.67 % of cases and was followed by colorectum and retroperitoneal sarcoma (16.67 %

each). In the young adult group, the 10 commonest cancers were colorectum (14.51%), breast (13.74%), skin+soft tissue (10.69%), testis (8.40%), gall bladder (7.63%), CNS (6.11%), hepatocellular carcinoma (4.58%), oral cavity (4.58%), bone (3.05%), stomach (3.05%), lymphoma (3.05%) and cervix (3.05%). In the middle age group, the 10 commonest cancers were breast (24.89%), gall bladder (9.37%), colorectum (8.20%), stomach (6.15%), periampullary (6.00%), Penis (4.83%), cervix (4.39%), oral cavity (4.39%), skin+soft tissue (4.25%) and HCC (4.25%). In the geriatric age group, the 10 commonest cancers were gall bladder (13.93%), prostate (12.31%), breast (10.38%), colorectum (9.29%), stomach (8.74%), penis (7.10%), skin+soft tissue (5.74%), urinary bladder (4.64%), periampullary (4.37%) and esophagus (4.37%).

Table 4: Age wise distribution of 10 commonest cancers (n = 1186).

Pediatric			Young adult			Middle			Geriatric		
Site	No.	%	Site	No.	%	Site	No.	%	Site	No.	%
Kidney	4	66.66	Colorectal	19	14.51	Breast	170	24.89	Gall Bladder	51	13.93
Colorectal	1	16.67	Breast	18	13.74	Gall Bladder	64	9.37	Prostate	45	12.31
Retroperitoneal	1	16.67	Skin	14	10.69	Colorectal	56	8.20	Breast	38	10.38
			Testis	11	8.40	Stomach	42	6.15	Colorectal	34	9.29
			Gall Bladder	10	7.63	Periampullary	41	6.00	Stomach	32	8.74
			CNS	8	6.11	Penis	33	4.83	Penis	26	7.10
			Hepatocellular	6	4.58	Cervix	30	4.39	Skin	21	5.74
			Oral cavity	6	4.58	Oral cavity	30	4.39	Urinary bladder	17	4.64
			Bone	4	3.05	Skin	29	4.25	Periampullary	16	4.37
			Stomach	4	3.05	Hepatocellular	29	4.25	Esophagus	16	4.37
			Lymphoma	4	3.05						
			Cervix	4	3.05						
10 Commonest	6	100	10 commonest	108	82.44	10 commonest	524	76.72	10 commonest	296	80.87
Others	0	0	Others	23	17.56	Others	159	23.28	Others	70	19.13
Total	6	100	Total	131	100	Total	683	100	Total	366	100

Table 5: Cancers with maximum number and proportion of metastasis at presentation (n = 1186).

Site	Total no. of cases	Presented with metastasis	Percentage
Gall bladder	125	58	46.40%
Cervix	39	24	61.53%
Liver 2 ^{ndaries} , primary unknown	21	21	100.00%
Breast	226	19	8.41%
Colorectum	110	17	15.45%
Stomach	78	10	12.82%
Prostate	61	8	13.11%
Ovary	19	8	42.10%
Testis	18	8	44.44%
Hepatocellular carcinoma	46	8	17.39%
10 commonest	743	181	24.36%
Others	443	37	8.35%
Total	1186	218	18.38%

Table 6: Distribution of mortality of patients according to age and gender (n = 91).

Age (years)	Number	%
0-14	0	0.00
15-34	10	10.99
35-64	49	53.85
65+	32	35.16
Total	91	100.00
Gender	Number	%
Male	56	61.54
Female	35	38.46
Total	91	100.00

It is evident from Table 5 that 218 i.e. 18.38% of patients had distant metastasis at the initial presentation. 21 patients presented with liver secondaries without detected primary lesion. 58 out of 125 i.e. 46.40% of patients of cancer gall bladder presented with metastasis making it

the cancer with the highest no. of cases with mets at the initial presentation.

It is evident from Table 6 that maximum deaths occurred in the middle age group (53.85%) followed by geriatric age group (35.16%) and young adults (10.99 %). There was no death among the pediatric patients. 61.54% of the deaths occurred among male patients and 38.46 % among female patients.

It is evident from Table 7 that the 10 commonest causes of deaths were gall bladder (14.29%), colorectum (13.17%), stomach (9.88%), and hepatocellular carcinoma (7.68%), CNS (brain + spinal cord) (5.49%), periampullary (5.49%), esophagus (4.40%), prostate (4.40%), oral cavity (4.40%) and skin+soft tissue (4.40%).

Table 7: Distribution of 10 commonest causes of mortality (n = 91).

Site of cancer	Total	
	Number	Percent
Gall bladder	13	14.29
Colorectum	12	13.17
Stomach	9	9.88
Hepatocellular carcinoma	7	7.68
CNS (brain+spinal cord)	5	5.49
Periampullary	5	5.49
Esophagus	4	4.40
Prostate	4	4.40
Oral cavity	4	4.40
Skin+soft tissue	4	4.40
10 commonest	67	73.60
Others	24	26.40
Total	91	100.00

Figure 1: Time trend of cancer admissions from 2010 to 2014.

During 2010-2014, 39213 patients were admitted in surgical wards out of which 1186 patients were of cancer which accounts for 3.02% of total admissions in surgical wards. The proportion of cancer patients increased from 2.45% from 2010 to 4.27% in 2014.

DISCUSSION

With the advent of the era of genetics, cancer has become one of the most researched diseases of the modern times and its etiopathogenesis is no more a mystery. Epidemiological studies are a major part of the research on cancer. The NCRP (National Cancer Registry Programme) with its pan-India network maintains a National database of cancer based on both population as well as hospital based registries and publishes the latest trends of cancer epidemiology. WHO (World Health Organization) publishes global as well as individual National data on cancer.

The incidence and mortality of various cancers is known to have age and gender predisposition. Various geographical areas, too, have different patterns of cancer distribution based on the genetic structure, socio-cultural

factors, and levels of pollution, life-style and exposure to carcinogens of the local residents.

This variation can be as much as 100 folds in certain cases. A systematic study of the distribution of cancer incidence, prevalence and mortality helps in charting out a trend of cancer over time and formulating important preventive strategies e.g. long time observation of relation of smoking to lung cancer led to a massive anti-smoking campaign and decline in incidence as well as mortality due to lung cancer.¹⁴

Incidence

In the present study, cancer accounted for 3.02% of the total admissions in surgical ward between 2010-2014. Globally, 14 million new cases of cancer are diagnosed each year.³ The age standardized cancer incidence rate is 182.3 per 100,000 population.⁷ In India, the annual incidence of cancer in India is 10.15 lakhs. The age standardized incidence rate is 94.0 per 100,000 population.^{8,9}

The proportion of cancer patients in present study is high because it is a hospital-based study and there is no cancer hospital in the Vindhya region, hence all cancer patients seek initial consultation in S. S. Medical College before being referred to specialized cancer hospitals. In order to estimate the prevalence of cancer in population, a population-based study is required.

Time trend of cancer - In the present study, the proportion of cancer patients has increased significantly from 2.45% to 4.27% from 2010 to 2014. Globally, while cancer rates are decreasing in the Western world, they are increasing in less developed and economically transitioning countries, including Eastern Europe, because of adoption of unhealthy western lifestyles such as smoking and physical inactivity and consumption of calorie-dense food. Cancers once known as diseases of industrialized countries e.g. lung, colon, and breast cancers, now commonly occur in economically transitioning and less developed countries. Most of these countries are also disproportionately affected by cancers related to infectious agents e.g. cervix, liver, and stomach cancers which are potentially preventable. The share of cancer burden in less developed countries is projected to increase from 56% of the world total in 2008 to >60% in 2030 because of increasing trends in cancer rates, expected increase in life expectancy and growth of population.¹⁰ The incidence of cancer is expected to increase from 14 million to 22 million cases per year and cancer mortality from 8.2 million to 13 million per year over next 2 decades.³

In India, cancer is on rise. Among males, except the Mumbai PBCR, all HBCRs and PBCRs have noted an overall increase in the AAR (age adjusted incidence rates) for cancer with Bangalore PBCR measuring the highest APC (annual percentage change) of 2.69. Among

the newer PBCRs, where statistics from 2003-2011 was calculated, except Mizoram and Sikkim states, all PBCRs and HBCRs saw an upward trend with Thiruvananthapuram PBCR recording the highest APC of 2.75. Among females, among the older PBCRs and HBCRs, except Delhi and Mumbai, the other ones showed a trend of increase in the overall incidence of cancer with Bangalore PBCR recording an APC of 2.90. In the newer PBCRs, except the Sikkim state, Mizoram state, Imphal West Dist and Dibrugarh Dist registries which showed a downward trend, the other registries showed an increasing trend with Kamrup Urban Dist measuring 11.38% and Thiruvananthapuram PBCR an APC of 3.48%.

A finding in the current study are in coherence with National data and is attributed to changing life-style and increasing pollution leading to accumulation of carcinogens and build-up of mutations as well as significant increase in the diagnosis of occult cancers with the help of advanced diagnostic modalities.

Incidence according to age

Various cancers show various patterns of incidence in different age groups. The commonest relation with age is a progressive increase in incidence from near zero in childhood and adolescence to a high rate in old age e.g. Cancers of skin, lung, gastrointestinal and urinary tracts where annual incidence may be 1000 times greater above 75 than before 25. Some cancers are typically found in pediatric population e.g. Retinoblastoma and nephroblastoma. Many other cancers have bimodal or multimodal distribution e.g. Breast cancer and lymphomas. In the present study, 57.59% patients were of middle age group, 30.86% were of geriatric age group, 11.05% young adults and 0.50% of pediatric age. Globally, life-time risk of developing cancer before the age of 75 years is 18.5%. (males 21.0%, females 16.4%).⁷ In India, life-time risk of developing cancer before the age of 75 years in India is 10.1% (males 10.2 %, females 10.1%).⁸

Most cases of cancer are seen in the age group of 35-64 years. Among males of this age group, incidence varies from 55.2 % in Chandigarh to 62.7% in Guwahati. Among females, it varies from 64.0% in Thiruvananthapuram to 74.3% in Dibrugarh. Among both the sexes, the figures vary from 60.3% in Chandigarh to 67.6% in Dibrugarh. The geriatric age group (65+ years) is the second commonest to be affected by cancer. Among males, the incidence in this age group varies from 19.2% in Mumbai to 30.1% in Guwahati. Among females, it varies from 14.2% in Mumbai to 20.0% in Thiruvananthapuram. Among both genders, it varies from 16.9% in Mumbai to 24.5% in Guwahati. Next are the young adults (15-34 years) with incidence among males varying from 5.6% in Guwahati to 13.3% in Mumbai. Among females, it varies from 8.7% in Guwahati to 11.9% in Thiruvananthapuram. Among both genders, it

varies from 6.8% in Guwahati to 12.5% in Mumbai. Pediatric cancers form rest of the bulk with incidence varying from 1.7% in Guwahati to 10.9% in Chandigarh among males, from 1.8% in Chennai to 6.4% in Chandigarh among females and from 1.8% in Guwahati to 8.9% in Chandigarh among both sexes.⁹ Findings of the study are consistent with the National data as in almost all cancer registries, the middle aged people account for the highest number of cancer cases, followed by geriatric age group, young adults and children. Epidemiological studies on cancer give results close to accurate when the study is restricted to the middle age group individuals because pediatric and young adult groups have a relatively less incidence of cancer and incidence of cancer in geriatric age group is often underestimated due to lack of universal diagnosis and reporting.

Incidence according to gender

In the present study, 50.25% patients were male, 49.75% were female. Globally, the age standardized cancer incidence rate is 205.4 for males and for females, it is 165.3 per 100,000 population.⁷ In India, annual incidence of cancer is 4.77 lakh among males and 5.37 lakh among females. The age standardized incidence rate is 92.4 among males and among females, 97.4 per 100,000 population.^{8,9} Males constitute 51.3% varying from 46.2% in Bengaluru to 59.2 % in Guwahati and females constitute 48.7% varying from 40.8% in Guwahati to 53.8% in Bengaluru.⁹ Findings of the present study are corroborating with the National data. It varies in respect to the Global data. Globally, incidence of cancers of breast and cervix which constitute the major bulk of female cancers are on fall in most of the advanced nations because of effective preventive and treatment strategies while the cancers of lungs which are known to affect males more are still unanswered and are high on incidence making males suffer more. In India, cancers of breast and cervix both are on rise, making the females share a bigger burden of cancer as compared to the global data.

Proportion according to religion

In the present study, 95.78% patients were Hindus, 4.22% were Muslims. This reflects the normal demographic distribution of the local region. Rewa District has 95.93% Hindus and 3.61% Muslims.¹¹

Proportion according to socio-economic status

In the present study, 70.40% patients were BPL (below poverty line), rest 29.60% were APL (above poverty line). Compared with other common surgical diseases, the distribution of cancer is more favored towards socio-economically affluent individuals due to the lifestyle hazards they are subjected to.

Proportion according to residence

In the present study, 67.2% patients were from rural areas, rests 32.80% were from urban areas. In Rewa, urban population is 16.73%.¹¹ Distribution of cancer is more favored towards urban areas due to more exposure to risk factors.

Proportion according to addiction

25.97% patients were addicted to tobacco, 18.63% were addicted to alcohol and tobacco both while 1.52% admitted to taking alcohol alone. But the data does not

conclude about any particular correlation of addiction and cancer and to find out a definite correlation, a quantitative and temporal correlation between addiction and cancer must be studied for a sufficiently long duration.

Proportion according to diet

75.89% patients were vegetarians which reflect the common dietary pattern of the region. To find out a correlation of diet with cancer, a longer duration study with detailed history of exposure to particularly known carcinogenic food agents must be done.

Table 8: Incidence according to site.

Present study				In India ¹²				Globally ⁷			
Males		Females		Males		Females		Males		Females	
Site	%	Site	%	Site	%	Site	%	Site	%	Site	%
Colorectal	11.24	Breast	37.12	Lungs	11.3	Breast	27.0	Lungs	16.7	Breast	25.2
Prostate	10.24	Gall bladder	12.88	Oral	11.3	Cervix	22.9	Prostate	15.0	Colorectal	9.2
Penis	10.24	Colorectal	7.28	Stomach	9.1	Colorectal	5.1	Colorectal	10.0	Lungs	8.8
Gall bladder	8.22	Cervix	6.61	Colorectal	7.7	Ovary	5.0	Stomach	8.5	Cervix	7.9
Stomach	7.55	Stomach	5.59	Pharynx	6.6	Oral	4.3	Liver	7.5	Uterus	4.8
Skin	7.05	Cholangio	4.06	Esophagus	5.7	Stomach	3.7	Urinary bladder	4.4	Stomach	4.8
Periampullary	6.88	Skin	3.73	Larynx	4.8	Lungs	3.1	Esophagus	4.3	Ovary	3.6
Oral cavity	6.88	Periampullary	3.22	Leukaemia	4.1	Esophagus	2.7	Kidney	2.9	Thyroid	3.5
Hepatocellular	4.53	Hepatocellular	3.22	Prostate	4.0	Leukaemia	2.4	NHL	2.9	Liver	3.4
Urinary bladder	4.03	Ovary	3.22	Liver	3.6	Uterus	2.3	Leukaemias	2.7	NHL	2.5

Proportion according to stage of presentation

18.38% of patients had proven distant metastasis at initial presentation. A detailed investigation and work up would have detected many covert metastases.

Disparity with the National and the global data is because the present study has been conducted on patients admitted in Surgical ward and only surgically respectable or manageable patients were included in the study and many of the patients admitted opted for further treatment in cancer hospitals after being diagnosed here. Patients of cancer of cervix seek consultation in the department of gynecology and only those patients turn up in the department of surgery who have developed obstructive uropathy or malignant ascites, hence the low incidence of cancer cervix in the study. Similarly in case of males, oral cavity cancers is the commonest throughout India but most of the cases seek consultation in the department of ENT/dentistry and only those cases with certain surgical complications report in the department of surgery.

Mortality

In the present study, cancer accounted for 3.64% of all deaths in the surgical ward and the case fatality rate of

cancer was 7.67%. Globally, 8.2 million people die of cancer each year.³ The age standardized cancer mortality rate is 102.4 per 100,000 population.⁷ In India, the annual mortality of cancer in India is 6.83 lakhs. The overall mortality rate is 64.5 per 100,000 population.^{8,9}

The reason behind high proportion of deaths of cancer patients in the present study is that it is a hospital-based study and there is no cancer hospital in the Vindhya region, hence all cancer patients seek initial consultation in the S. S. medical college before being referred to specialized cancer hospitals. Many patients present in advanced stages and are unable to seek care in specialized cancer hospitals and die during palliative treatment in S. S. medical college, Rewa. In order to estimate mortality rate of cancer in population, a population-based study is required.

Mortality according to age

In the present study, maximum deaths occurred in the middle age group (53.85%) followed by geriatric age group (35.16%) and young adults (10.99%). There was no death among the pediatric patients. Globally, life-time risk of dying due to cancer before the age of 75 years is 10.4%. Among males, it is 12.7% and among females,

8.4%.⁷ In India, life-time risk of dying due to cancer before the age of 75 years in India is 7.1%. Among males, it is 7.8% and among females, 6.5%.⁸

The reason behind the relatively less proportion of total deaths attributed to cancer among the geriatric age group as compared to the global data is that a significant proportion of the deaths in this group remains uninvestigated and are attributed to non-malignant causes, due to lack of universal autopsy and mortality reporting system. A correct picture can be obtained if all patients in this age group are cared after properly and are diagnosed thoroughly and properly for their diseases and all deaths are subjected to autopsy to confirm the exact

cause of death and a National databank of the same is maintained as is done in developed nations.

Mortality according to gender

In the present study, 61.54% of deaths occurred among male patients and 38.46% among female patients. Globally, the age standardized cancer mortality rate for males is 126.3 and for females, 82.9 per 100,000 population. In India, the annual mortality rate of cancer in India is 3.57 lakh among males and 3.26 lakh among females. The overall mortality among males is 69.7 and among females, 60.2 per 100,000.^{8,9}

Table 9: Mortality according to site.^{7,12}

Present study				In India ¹²				Globally ⁷			
Males		Females		Males		Females		Males		Females	
Site	%	Site	%	Site	%	Site	%	Site	%	Site	%
Stomach	12.49	Gall bladder	20.00	Lung	13.7	Breast	21.5	Lung	23.6	Breast	14.7
Gall bladder	10.70	Colorectal	17.14	Stomach	11.4	Cervix	20.7	Liver	11.2	Lung	13.8
Colorectal	10.70	Cervix	8.57	Oral	10.2	Colorectal	6.4	Stomach	10.1	Colorectal	9.0
Prostate	7.14	Thyroid	8.57	Colorectal	7.8	Ovary	6.0	Colorectal	8.0	Cervix	7.5
Periampullary	7.14	Hepatocellular	8.57	Pharynx	7.6	Stomach	5.6	Prostate	6.6	Stomach	7.2
Hepatocellular	7.14	Stomach	5.71	Esophagus	7.1	Oral	4.8	Esophagus	6.0	Liver	6.3
Esophagus	5.36	Oral	5.71	Liver	4.7	Lungs	4.6	Leukemia	3.3	Ovary	4.3
CNS	5.36	CNS	5.71	Leukemia	4.5	Esophagus	4.1	Urinary bladder	2.6	NHL	2.4
Skin	5.36	Cholangio	5.71	Larynx	4.4	Leukemia	3.3	NHL	2.5	Uterus	2.1
Hematological	5.36	Periampullary	2.86	Prostate	3.4	Uterus	1.5	Kidney	2.0	Thyroid	0.8

Absences of death due to breast cancer in the present study can be explained by availability in the institution of various treatment modalities for the management of breast cancer including surgery, chemotherapy, radiotherapy and hormonal therapy which ensures a near complete treatment. Also many patients of breast cancer might have died due to other co-morbid conditions in separate admissions and hence not included in the study.

CONCLUSION

This was a 5 year retrospective+prospective hospital based study. A longer duration prospective population based study with direct interviewing of patients regarding various relevant traits and following their management on a long-term basis would be an ideal and more comprehensive study. Nonetheless, this study helps in pointing out the prevalent trends in cancer in the Vindhya region. The battle against cancer must be fought on many frontiers and a study to find out its determinants will act as a foot soldier and prepare the cancer management team for greater challenges.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the institutional ethics committee

REFERENCES

1. World Cancer Factsheet, IARC (International Agency for Research on Cancer), WHO, 2014. Available at http://publications.Mcancerresearchuk.org/downloads/Product/CS_REPORT_WORLD.pdf. Accessed on 21 January 2016.
2. Robbins Basic Pathology 8th Edition; 2007:175.
3. Latest World Cancer Statistics, Press Release No. 223. IARC (International Agency for Research on Cancer), WHO; 2013.
4. Cancer in India: prevalence, management and outcome prospects for research. Subrata Sinha Department of Biochemistry All India Institute of Medical Sciences New Delhi, 2004. Available at http://www.euindiacoop.org/2009/downloads/subrata_sinha.pdf. Accessed on 21 January 2016.
5. Mallath MK, Taylor DG, Badwe RA, Rath GK, Shanta V, Pramesh CS, et al. The growing burden of cancer in India: epidemiology and social context. *Lancet Oncology*. 2014;15(6):e205-212.

6. Cancer epidemiology, prevention and control, current science. 2004;86(4):25.
7. Globocan, World Fact Sheet, section of cancer information. International Agency for Research on Cancer, Lyon, France. 2013
8. Stewart BW, Kleihues P. According to WHO, World Cancer Report Ed. by; 2013.
9. The National Cancer Registry Programme Consolidated Reports 2007-2011, ICMR (Indian Council of Medical Research). Available at http://www.ncrpindia.org/ALL_NCRP_REPORTS/HBCR_REPORT_2007_2011/index.htm. Accessed on 21 January 2016.
10. Jemal A, Sentalis C. Global patterns of cancer incidence and mortality rates and trends, cancer epidemiology biomarkers and prevention. 2010;19(8):1893-907.
11. The Census of India, 2011. Available at https://en.wikipedia.org/wiki/2011_Census_of_India. Accessed on 21 January 2016.
12. Globocan, World Fact Sheet, section of cancer information. International Agency for Research on Cancer, Lyon, France. 2013.
13. Kapoor VK, MvMichael AJ. Gallbladder cancer: an Indian disease. *Natl Med J India*. 2003;16:209-13,
14. Trends in cancer incidence and mortality. Cancer surveys. Cold Spring Harbor Laboratory Press, New York, Doll R, Fraumeni J, Muir C, eds. 1994;(19-20).
15. Time trend of cancer incidence, 1982-2010, NCRP. Available at http://www.ncrpindia.org/Annual_Reports.aspx. Accessed on 21 January 2016.

Cite this article as: Shukla D, Patel VK, Chhari AS, Dubey CS, Garg RK, Singh AP, et al.

Epidemiological profile of cancer patients in surgery ward of a tertiary teaching hospital in the Vindhya region with special reference to high proportion of carcinoma gall bladder. *Int Surg J* 2016;3:2025-33.