# **Original Research Article**

DOI: http://dx.doi.org/10.18203/2349-2902.isj20183194

# Epidemiological analysis of 1637 cases of animal bite injuries

Mudassir M. Wani<sup>1\*</sup>, Zia-Ul-Sabah<sup>2</sup>

<sup>1</sup>Department of Surgery, Government Medical College, Srinagar, India

Received: 12 May 2018 Accepted: 26 June 2018

## \*Correspondence:

Dr. Mudassir Maqbool Wani, E-mail: drmudds@gmail.com

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## **ABSTRACT**

**Background:** This study was conducted prospectively over a period of two years in accident and emergency department of Government Medical College, Srinagar. The main objectives of the study included finding out epidemiological aspects of animal bites, categorising bites and analysing associated injuries.

**Methods:** From April 2009 to April 2011, more than 1800 patients presented to accident emergency department with injuries due to human animal conflict. Out of them, 1673 patients had only bite injuries and were treated on out patient basis were included in the study.

**Results:** Study revealed incidence in males (69.27%) being higher than in females. The age group most commonly involved was 0-10 years and rural population (60.72%) was more affected as compared to urban population. Furthermore, dogs were most common (88.46%) animals to be involved in bites. The commonest bite category was class 2 among all the bites (58.57%) while as type 1 was least common (9.74%). Injury pattern showed that most common site of bite was lower limbs, left leg (56.90%) being more common than the right.

**Conclusions:** We conclude from this study that animal bites usually involve age group of less than 10 years. Males and rural population are more commonly involved. Dog is most common animal involved in such conflicts. We found lower limbs are most common sites involved with left lower limb being more commonly bitten as compared to right.

**Keywords:** Animal bites, Bite category, Injury pattern, Rabies

# INTRODUCTION

The human animal interaction is ancient. Anthro-zoology (also known as human–non-human-animal studies or HAS) is the study of interaction between humans and other animals. One of the most important aspect of this branch is the quantifying of the positive effects of human-animal relationships on either party and the study of their interactions.<sup>1</sup>

Among those interactions the bond between dogs and humans is enduring. But question arises, why do animals, particularly pet ones, bite? Well it is a conundrum. Human animal conflict dates back to the prehistoric times. The nature of conflict may have not changed a lot but perception about the outcome has changed a lot from past to present, earlier concern may have been about injury only but now infections like rabies are a priority to both patient as well as medical practitioners.

Diseases like rabies still remain a challenge in developing and underdeveloped world particularly South East Asia, with an estimated 45% deaths worldwide due to rabies occurring in that part.<sup>2</sup> The condition in India being more worse with about 36% of rabies deaths being reported worldwide and showing no decline in incidence from last one decade.<sup>3,4</sup>

<sup>&</sup>lt;sup>2</sup>Department of Medicine, King Khalid University, Abha, Aseer, Saudi Arabia

This present study was carried in Northern State of India with a very high incidence of human - dog and human-wildlife conflicts.<sup>5</sup> With this background and scarcity of literature available regarding animal bites pattern, we carried out this study.

The study was conducted in one of the main tertiary care referral hospital of the state, having well established anti rabies clinic. The main purpose of study was finding out epidemiological aspects of animal bites, categorizing and analyzing bite associated injuries.

#### **METHODS**

This study was carried out in Accident Emergency Department of Medical college affiliated hospital. It is one of the main referral centres of the state having a well-established Anti Rabies Clinic. Study was carried from April 2009 to April 2011 and 1673 patients were included.

## Inclusion criteria

- Patients who reported primarily to our hospital.
- Patients who were treated on Out-patient basis after primary management.
- Patients who provided detailed history.

#### Exclusion criteria

- Patients who were referred to our hospital after receiving primary treatment
- Patients who needed admission in hospital for more than 24 hours
- Patients who needed any operative surgery due to bite or bite related injury.

#### **RESULTS**

A total number of 1673 patients were included in study. All injuries were classified as per WHO Classification, Table 1.

Table 1: WHO classification of Animal Bites.

| Class | Definition                              |
|-------|---|
| I.    | Touching or feeding of animals.         |
|       | Licks on intact skin                    |
| II.   | Nibbling of uncovered skin.             |
|       | Minor scratches or abrasions without    |
|       | bleeding.                               |
|       | Licks on broken skin                    |
| III.  | Single or multiple transdermal bites or |
|       | scratches.                              |
|       | Contamination of mucous membrane        |
|       | with saliva (i.e. licks)                |

The number of male patients included 1134(69.27%) as compared to females 503(30.72%). Out of 1673, 994 (60.72%) patients belonged to rural areas while as 643(39.27%) patients were from urban areas, Table 2.

Table 2: Sex and Area distribution.

| Distribution      | Characteristics |              |
|-------------------|-----------------|--------------|
| Sex distribution  | Male            | Female       |
|                   | 1134 (69.27%)   | 503 (30.72%) |
| Area distribution | Rural           | Urban        |
|                   | 994 (60.72%)    | 643 (39.27%) |

All patients were enquired about the animal involved. The most common animals implicated in bite cases were dogs, 1468, 89.67% Table 3.

Table: 3 Animals involved with corresponding age group distribution.

| Age group        | Dog          | Bear      | Cat       | Horse     | Monkey    | Other      | Total |
|------------------|--------------|-----------|-----------|-----------|-----------|------------|-------|
| 0-10             | 496          | 7         | 0         | 0         | 3         | 21         | 528   |
| 11-20            | 279          | 2         | 2         | 1         | 2         | 27         | 312   |
| 21-30            | 185          | 2         | 0         | 2         | 1         | 14         | 204   |
| 31-40            | 234          | 2         | 2         | 3         | 2         | 26         | 269   |
| >40              | 274          | 1         | 9         | 7         | 2         | 31         | 324   |
| Total percentage | 1468 (89.67) | 14 (0.08) | 13 (0.07) | 13 (0.07) | 10 (0.06) | 119 (7.20) | 1637  |

Table 4: Age group and bite category.

| Age group | Class 1   | Class 2    | Class 3    |              |
|-----------|-----------|------------|------------|--------------|
| 0-10      | 48        | 328        | 152        | 528 (31.55%) |
| 11-20     | 35        | 180        | 97         | 312 (18.64%) |
| 21-30     | 26        | 131        | 47         | 204 (12.19%) |
| 31-40     | 34        | 156        | 79         | 269 (16.07%) |
| >40       | 20        | 185        | 119        | 324 (19.36%) |
|           | 163(9.74) | 980(58.57) | 494(29.52) | 1673         |

The patients were divided into 4 age groups: 0 to 10,11 to 20,31 to 40 and more than 40 years. Highest incidence animal bites of 31.55 % was found in age group of 0-10 years, followed by age groups of >40 years and 11-20 having incidence of 19.36% and 18.64% respectively. Least common age group was 21-30 years having incidence of 12.19%, Table 4. Out of 1637 cases many

patients had multiple bites. As such total number of bite injuries in these patients were about 1685. Lower limb was the most common site of bites constituting about 65.28% cases. Besides left lower limb was more commonly involved as compared to right, 57% vs 42%, Table 5.

Table 5: Site of involvement in bites.

| Site                    | Total number  |                          | Most common age group involved |
|-------------------------|---------------|--------------------------|--------------------------------|
| Head and neck           | 73 (04.30%)   |                          | 0-10 years (47)                |
|                         |               | Right Limb: 141 (52.61%) |                                |
| Upper limb              | 268 (15.90%)  | Left Limb: 109 (40.67%)  | 0-10 years (84)                |
|                         |               | Both Limbs:18 (6.71%)    |                                |
| Lower limb              |               | Right Limb: 464 (42.18%) |                                |
|                         | 1100 (65.28%) | Left Limb: 626 (56.90%)  | 0-10 years (327)               |
|                         |               | Both Limbs: 10 (0.9%)    |                                |
| Trunk and other regions | 244 (14.48%)  |                          | 0-10 years (103)               |

The treatment part included thorough washing of wound, active immunization using cell derived vaccines. In category 3, local infiltration using ARS was also done. All patients received first dose free. Immunoglobulin had to be bought by patients on their own. If patients could not afford , same was provided by social organisation (Help Poor Foundation). For further doses patients were referred to different outlets of Anti Rabies Clinic as per area patient belonged.

# **DISCUSSION**

About half of Americans will be bitten by an animal or another human being during their lifetimes, as such animal bites account for 1% of all emergency department visits.<sup>6</sup> It is estimated that 28% of all children have a calculated risk of being bitten by a dog at least once.<sup>7</sup> The study undertaken presently focused on epidemiological aspects of animal bites, categorizing and analyzing animal bite associated injuries. We could find good number of publications about rabies as well as bite associated infections but could not find much written about animal bite injury pattern and categorization of same.

The study revealed that rural population and males in particular had higher incidence in terms of total number of bites. This is in accordance with previous studies. 8-10 In addition we found that stray dogs were the main biting animals, similarly to other studies. 8-11 But in contrast to majority of other studies we found bear bites second in frequency rather than cat.

The main reason for the same can be explained on the basis that in the area of Jammu and Kashmir, as per world

life SOS, there are highest number of human conflicts with Asiatic Black bears and leopards. We found that the most common age group being bitten by dogs were less than 10 years of age. The percentage being about 31.55%, which is less as seen in some studies in which 57% patients belonged to same group. 12-13 The difference can be understood from the fact that the previous studies have been conducted in developed countries where having pets is common. On another hand, our study has been carried out in a place where having pets is not common as it is often considered against social and religious customs. The second common age group bitten belonged to people aged more than 40, n= 324(19.36%) cases. The explanation can be given that after 40 years the strength is decreased and running away from animal may not be easy.

We estimated class of bite injuries. We found that Class 2 injuries being most common, 58.57%. But we feel Class 1 injuries were underestimated as minor cases are either treated locally or managed by primary care physicians, same has been noticed in other studies.<sup>7</sup> Finding exact incidence of same was beyond our study.

The final aspect of study was to analyse injury pattern in animal bites. We found that some patients had more than one bite, as such we had more bite injuries as compared to number of patients. The lower limb injuries were most common, n=1100. In lower limb, left limb bites 626 (56.90%) were much more than right limb 464 (42.18%). The reason for the same is that at the time of animal attack people try to run away using right leg first and as such left leg is often bitten.

In 10 patients both limbs were bitten. Upper limbs were involved next in frequency of bites; right limb bitten in

141 (52.61%); left Limb in109 (40.67%) and both limbs in18(6.71%). The results for upper limb are somewhat similar to other studies[14]. Trunk and abdomen injuries ranked third n=244(14.48%) while as head and neck region had minimum bites, n=73 (04.30%).

In contrast to other studies in India, we found that majority of patients we received had kept wound open (No Covering Dressing) and usually had applied some antiseptic. Although we also found that knowledge about wound washing was lacking in majority.<sup>4</sup>

#### **CONCLUSION**

We conclude from this study that animal bites usually involve age group of less than 10 years. Males and rural population are more commonly involved. Category 2 bites are most common ones seeking medical attention. Injury pattern revealed that lower limbs were the most commonly involved and left lower limb being more commonly bitten as compared to right.

Funding: No funding sources Conflict of interest: None declared

Ethical approval: The study was approved by the

Institutional Ethics Committee

## **REFERENCES**

- 1. Mills, Daniel S. Anthrozoology. The Encyclopedia of Applied Animal Behaviour and Welfare. CABI. 2010:28-30.
- 2. Gongal G, Wright AE. Human rabies in the WHO Southeast Asia Region: forward steps for elimination. Adv Prev Med. 2011;2011:383870.
- 3. Dutta JK. Human rabies in India: epidemiological features, management and current methods of prevention. Trop Doct. 1999;29:196-201.
- 4. Sudarshan MK, Madhusudana SN, Mahendra BJ, Rao NS, Ashwath Narayana DH, Abdul Rahman S, et al. Assessing the burden of human rabies in India:

- results of a national multi-center epidemiological survey. Int J Infect Dis. 2007;11:29-35
- 5. Reducing Conflict, Wild life SOS of India. Available at. www.wildlifesos.org.
- 6. Griego RD, Rosen T, Orengo IF, Wolf JE. Dog, cat, and human bites: a review. J Am Acad Dermatol. 1995;33(6):1019-29.
- 7. George K, Adesiyun A. An investigation into the prevalence of dog bites to primary school children in Trinidad. BMC Public Health. 2008;8:85.
- 8. Sudarshan MK, Mahendra BJ and Ashwathnarayan DH. A community survey of dog bites anti-rabies treatment, rabies and dog population management in Bangalore City. J Com Dis. 2001;33(4):245-51.
- 9. Singh Jagvir, Jain DC, Bhatia Rajesh et al. Epidemiological characteristics of rabies in Delhi and surrounding areas, 1998, Indian Paediatr. 2001;38:1354-60.
- 10. Agarwal N, Reddaiah VP. Knowledge, attitude and practice following dog bite: A community based epidemiological study. Health Population. Perspect Issues. 2003;26(4):154-61.
- 11. National Institute of Communicable Diseases, Rabies: A major public health problem. CD Alert. 2000;4(10)
- 12. Benson LS, Edwards SL, Schiff AP, Williams CS, Visotsky JL. Dog and Cat bites of hand: Treatment and cost assessment. J Hand Surg. 2006;31A:468-73.
- 13. Sacks JJ, Lockwood R, Hornreicht J, Sattin RW. et al. Fatal dog attacks,1989-1994. Peadiatrics. 1996;97:891-5.
- 14. Marianne NW, Lars D. Dog bite injuries to hand. J Plas Surg Hand Surg. 2011;455(2):96-101.

**Cite this article as:** Wani MM, Sabah ZU. Epidemiological analysis of 1637 cases of animal bite injuries. Int Surg J 2018;5:2799-802.