

Original Research Article

Predictive factors for intraoperative excessive bleeding in multinodular goitre with or without hyperthyroidism

Ramesh C. Sagar, Gudekote Ravikumara*, V. B. Malipatil

Department of General Surgery, Raichur Institute of Medical Sciences, Raichur, Karnataka, India

Received: 16 July 2018

Revised: 19 September 2018

Accepted: 27 September 2018

*Correspondence:

Dr. Gudekote Ravikumara,

E-mail: ravikumar.gudekote@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Multinodular goitre disease frequently results in and is the most common cause of hyperthyroidism. It also often results in an enlarged thyroid. It also known as toxic diffuse goiter, is an autoimmune disease that affects the thyroid. This study evaluates all the factors that cause intra-operative blood loss and how it affects the Grave's disease.

Methods: This study was conducted in Raichur Institute of Medical Sciences Raichur on 200 patients with multinodular goitre disease, who underwent thyroidectomy during the period between November 2010 to July 2015.

Results: The majority of patients were females which constitute about 76.3% with a median age of 33 years. The median period between the onset of the disease and operation was 15 months. Weight of thyroid in grams was 40. Post-operative hospital stay was 4 hours. Univariate analysis revealed that the strongest correlation of AIOBL was noted with the weight of thyroid ($p < 0.001$). Additionally, AIOBL was correlated positively with the period between disease onset and surgery ($p < 0.001$) and negatively with preoperative free T4 ($p < 0.01$). Occurrences of postoperative complications, such as recurrent laryngeal nerve palsy or hypoparathyroidism, and postoperative hospital stay were not correlated with AIOBL.

Conclusions: For multinodular goitre, for excessive bleeding during surgery, a large goiter presented as a predictive factor, and transfusion of blood should be considered in cases in which goitre weighs more than 200 g.

Keywords: Intra-operative excessive bleeding, Multinodular goitre,

INTRODUCTION

Multinodular goitre, also known as toxic diffuse goiter, is an autoimmune disease that affects the thyroid.¹ It frequently results in and is the most common cause of hyperthyroidism.² It also often results in an enlarged thyroid. Signs and symptoms of hyperthyroidism may include irritability, muscle weakness, sleeping problems, a fast heartbeat, poor tolerance of heat, diarrhea, and unintentional weight loss. Other symptoms may include thickening of the skin on the shins, known as pretibial

myxoedema, and eye bulging, a condition caused by Graves' ophthalmopathy. About 25 to 80% of people with the condition develop eye problems.³ The exact cause is unclear; however, it is believed to involve a combination of genetic and environmental factors.⁴ A person is more likely to be affected if they have a family member with the disease. If one twin is affected, a 30% chance exists that the other twin will also have the disease.⁵ The onset of disease may be triggered by stress, infection, or giving birth. Those with other autoimmune diseases such as type 1 diabetes and rheumatoid arthritis are more likely to be

affected. Smoking increases the risk of disease and may worsen eye problems. The disorder results from an antibody, called thyroid-stimulating immunoglobulin (TSI), that has a similar effect to thyroid stimulating hormone (TSH). These TSI antibodies cause the thyroid gland to produce excess thyroid hormone. The diagnosis may be suspected based on symptoms and confirmed with blood tests and radioiodine uptake. Typically, blood tests show a raised T₃ and T₄, low TSH, increased radioiodine uptake in all areas of the thyroid, and TSI antibodies. The three treatment options are radioiodine therapy, medications, and thyroid surgery. Radioiodine therapy involves taking iodine-131 by mouth, which is then concentrated in the thyroid and destroys it over weeks to months. The resulting hypothyroidism is treated with synthetic thyroid hormone. Medications such as beta blockers may control some of the symptoms, and antithyroid medications such as methimazole may temporarily help people while other treatments are having effect. Surgery to remove the thyroid is another option. Eye problems may require additional treatments. Graves' disease will develop in about 0.5% of males and 3% of females. It occurs about 7.5 times more often in women than in men.¹ Often, it starts between the ages of 40 and 60, but can begin at any age. It is the most common cause of hyperthyroidism in the United States (about 50 to 80% of cases). The exact cause is unclear; however, it is believed to involve a combination of genetic and environmental factors. While a theoretical mechanism occurs by which stress could cause an aggravation of the autoimmune response that leads to goitre disease, more robust clinical data are needed for a firm conclusion. Thyroidectomy is considered as a prompt therapy for multinodular goitre disease, but it has few disadvantages such as scar post operatively, need for hospitalization, operation complications, recurrent laryngeal nerve, hypoparathyroidism, and intra-operative bleeding. This study evaluates all the factors that cause intra-operative blood loss and how it affects the disease.

METHODS

This study was conducted on 200 patients with multinodular goitre disease, who underwent thyroidectomy during the period between November 2010 to July 2015.

The excluded patients were those who had undergone neck surgery before, those patients who had recurrent goiter, vocal cord paralysis, thyroid cancer requiring lymph node dissections. For a period of 14 days, all patients were administered with potassium iodide before the surgery. Conventional open surgery or neck surgery video assisted were performed. Closed suction drainage tubes before wound closure in all patients were given and removed them 1 to 3 days later. If voice change was observed, recurrent laryngeal nerve injury was recorded.

Those patients who had symptomatic hypocalcemia were given calcium supplements and vitamin D analogue.

Patients who within 12 months after surgery, who could discontinue the supplements and who could achieve normocalcemia were considered as transient hypocalcemia. Patients who had to continue the supplements for more than 12 months and who had < normal serum parathyroid hormone level were considered as permanent hypoparathyroidism. Variables of clinical were measured using AIOBL. To test the correlation between continuous variables and AIOBL, univariate analysis was performed. Students test, and regression analysis was used to compare between groups. A p value of less than 0.05 was said to be statistically significant. For multivariate analysis, P value of ≤ 0.10 was considered to be stepwise regression analysis to determine the risk factors for intra-operative bleeding.

RESULTS

The majority of patients were females which constitute about 76.3% with a median age of 33 years. The median period between the onset of the disease and operation was 15 months. Weight of thyroid in grams was 40. Post-operative hospital stay was 4 hours.

Table 1: Demographics of patients in the study and operative variables.

Demographics	
Mean age in years	33 (10-73)
Sex, male:female	50:150
Period between disease onset and surgery	15 (1-460)

Table 1 shows that the mean age in years of patients were 33, which ranged between 10 to 73 years, male to female ratio was 50:150. The period between disease onset and surgery was 15.

Table 2: Background of patients in the study and operative variables.

Indication for surgery	
Intolerance to antithyroid drug	80 (40%)
Patient preference	45 (22.5%)
Uncontrollable disease	45 (22.5%)
Large goiter with pressure symptoms	40 (20%)
Poor adherence	15 (7.5%)
Grave's ophthalmology	10 (5%)
Planned pregnancy	5 (2.5%)
Malignancy suspicion	1 (0.5%)
Thyrotoxic crisis	1 (0.5%)
Young age	1 (0.5%)
Failure of radioiodine therapy	1 (0.5%)
Pre-operative TSH at initial consultation (ng/mL)	0.004 (0.0-77.5)
Pre-operative free T4 at initial consultation (ng/mL)	2.5 (0.2-21.2)
Pre-operative steroid	35(17.5%)
Pre-operative β -blocker	80(40%)

Table 2 shows that intolerance to antithyroid drug and pre-operative β -blocker was the highest indication for surgery constituting 40% (80). The least was pre-operative TSH at initial consultation (ng/mL) which constituted to 0.0-77.5 (0.004).

Table 3: Variables of the surgery.

Variables of surgery	
Duration of surgery (mins)	185 (68-378)
Amount of blood loss (mL)	95 (10-1385)
Blood transfusion	5

Table 3 shows that duration of surgery in minutes was 185, amount of blood loss was 95 mL and 5 blood transfusions happened.

Table 4: Operative methods and surgical complications.

Operative methods	
Open surgery:video assisted neck surgery	120:70
Subtotal:near total/total thyroidectomy	135:55
Surgical complications	
Transient hypocalcemia	38 (19%)
Permanent hypoparathyroidism	1 (0.5%)
Transient recurrent laryngeal nerve palsy	5 (2.5%)
Permanent recurrent laryngeal nerve palsy	0 (0.0%)

Table 4 shows that open surgery:video assisted neck surgery was 120:70, subtotal:total/near total thyroidectomy was 135:55. Surgical complications were seen in 38 patients who had transient hypocalcemia and the least complications were seen in permanent recurrent laryngeal nerve palsy i.e. 0.

Table 5: Correlation between clinical variables and intra-operative blood loss (univariate analysis).

Continuous variables (regression analysis)	Correlation coefficient	P-value
Age in years	0.07	0.25
BMI	0.04	0.45
Period between disease onset and surgery	0.34	<0.001
Free T4 at initial consultation	0.20	<0.05
Weight of thyroid	0.70	<0.0001
Categorical variables (students t test)		
Sex		0.50
Operative methods		0.10
Type of surgical resection		0.50

Table 5 shows that regression analysis was done for age in years with correlation coefficient of 0.07, P-value of 0.25, BMI with correlation coefficient of 0.04, P value of 0.45, Period between disease onset and surgery with correlation coefficient of 0.34, P value of <0.001, Free T4

at initial consultation with correlation coefficient of 0.20, P value of <0.05 and weight of thyroid with correlation coefficient of 0.70, P-value of <0.0001.

DISCUSSION

Yamanouchi K et al reported that a total of 197 patients underwent thyroidectomy for multinodular goitre disease between 2002 and 2012.⁶ It was evaluated that clinical factors that would be potentially related to AIOBL retrospectively. The median period between disease onset and surgery was 16 months (range: 1e480 months). Conventional surgery was performed in 125 patients, whereas video-assisted surgery was performed in 72 patients.

Subtotal and near-total/total thyroidectomies were performed in 137 patients and 60 patients, respectively. The median weight of the thyroid was 45 g (range: 7.3e480.0 g). Univariate analysis revealed that the strongest correlation of AIOBL was noted with the weight of thyroid ($p < 0.001$). Additionally, AIOBL was correlated positively with the period between disease onset and surgery ($p < 0.001$) and negatively with preoperative free T4 ($p < 0.01$). Multivariate analysis showed that only the weight of the thyroid was independently correlated with AIOBL ($p < 0.001$). Four patients (2.0%) needed blood transfusion, including two requiring autotransfusion, whose thyroids were all weighing in excess of 200 g.

The amount of drainage during the initial 6 hours and days until drain removal was correlated positively with AIOBL ($p < 0.001$, each). Occurrences of postoperative complications, such as recurrent laryngeal nerve palsy or hypoparathyroidism, and postoperative hospital stay were not correlated with AIOBL. Grodski S et al conducted a study in which the most frequent absolute indication was the presence of a large goiter ($n = 8$; 13%) or associated thyroid nodule ($n = 6$; 10%).⁷

Ophthalmopathy, a relative indication, comprised the largest single group overall ($n = 18$; 29%); however, a significant number of patients ($n = 17$; 27%) elected surgery in the absence of a recognized indication. There was strong concordance (73%) between the recorded indication and the patients' survey response. Overall, there was a high level of satisfaction with surgery with 88% of respondents giving a satisfaction score of 7 or greater on a visual analog scale (VAS) (0-10). Erbil Y et al reported that the mean blood flow, MVD, CD-34 expression, and blood loss in group 1 patients were significantly lower than those in group 2 patients.⁸

There was a negative correlation between Lugol solution treatment and blood flow ($r(s) = -0.629$; $P = 0.0001$), blood loss ($r(s) = -0.621$; $P = 0.0001$), MVD ($r(s) = -0.865$; $P = 0.0001$), and CD-34 expression ($r(s) = -0.865$; $P = 0.0001$). According to logistic regression analysis, Lugol solution treatment resulted in a 9.33-fold decreased

rate of intraoperative blood loss. Karamanakos SN et al observed that a total, near-total and subtotal thyroidectomy was performed in 1,149,777 and 117 patients, respectively.⁹ Transient RLNP occurred in 34 (1.6%) and permanent in 19 (0.9%) patients. Multivariate logistic regression analysis showed that extended resection (OR-odds ratio-1.6), Graves' disease (OR 2.7), thyroiditis (OR 2.1), recurrent goiter (OR 2.3) and thyroid malignancy (OR 1.7) were all independent risk factors for transient RLNP, whereas Graves' disease (OR 2.2) and recurrent goiter (OR 1.7) emerged as independent risk factors for permanent RLNP. The rates of transient and permanent hypoparathyroidism were 27.8% and 4.8%, respectively.

Multivariate analysis for transient hypoparathyroidism revealed that the extent of surgical resection (OR 2.2), Graves' disease (OR 2.1), recurrent goiter (OR 1.7), female gender (OR 1.5) and specimen weight (OR 1.6) were independent predictors. However, the extent of surgical resection (OR 2.7), Graves' disease (OR 1.8), recurrent goiter (OR 1.5) and malignant disease (OR 1.5) were independent risk factors for permanent hypoparathyroidism. Postoperative wound infection and hematoma occurred in 6 (0.3%) and 27 (1.3%) patients, respectively. No correlation was observed between wound infection or postoperative hemorrhage and the extent of surgery.

Allannic H et al performed a prospective randomized study was performed in patients with hyperthyroid Graves' disease (GD) in order to compare long (18 months) and short term (6 months) antithyroid drug treatment on the remission rate.¹⁰ A therapeutic protocol was offered to all GD patients who had not been treated for this disease previously. All patients studied who followed the protocol were rechecked 2 yr after treatment was withdrawn, or earlier in the case of relapse. Of the patients having undergone long term treatment, 61.8% still were in remission 2 yr after treatment withdrawal, whereas only 41.7% of the patients treated for 6 months were in remission (P less than 0.05). Such findings clearly establish that treatment duration has a direct beneficial incidence on the remission rate. These results were confirmed by the fact that treatment for 18 months resulted in remission in 7 of 15 patients who had previously relapsed after a 6-month course of therapy. This improvement in relation to treatment duration might be due to the immunosuppressive action of carbimazole. No significant difference was observed between relapse and remission groups, regardless of treatment duration, for HLA ABDr, serum T3 and T4, and T3/T4 ratio determined before treatment. Only the thyroid-stimulating antibody levels determined at the time of diagnosis and at the end of treatment were higher in the relapse group, a difference that was relevant only globally, due to value scattering.

Thomusch O et al conducted a prospective study from 45 East German hospitals involving 7266 patients with

surgery for benign goiter.¹¹ High-volume providers (>150 operations per year) performed 69% (5042/7266), intermediate-volume providers 27% (50-150), and low-volume providers 4% (258/7266) of operations. The pattern of thyroid disease did not vary significantly, but there was a trend that small-volume providers tended to perform more operations for uninodular goiter and high-volume providers treated more patients with Graves' disease and recurrent goiter among the hospital groups. Extent of resection ($p < 0.0001$) and remnant size (multinodular goiter and recurrent goiter, $p < 0.001$), differed significantly, with total thyroidectomy being performed more often in hospitals with more than 150 operations compared to hospitals with an operative volume of less than 150 procedures per year. Despite the larger extent of resection and smaller remnant size, rates of recurrent laryngeal nerve (RLN) palsy or hypoparathyroidism were not increased. When the logistic regression analyses were fitted to evaluate the impact of risk factors on transient and permanent RLN palsy and hypoparathyroidism, larger extent of resection [relative risk (RR) 1.5-2.1] and recurrent goiter (RR 1.8-3.4) consistently evolved as independent risk factors. With hypoparathyroidism, additional significant factors included patient gender (RR 2.1-2.4), hospital operative volume (RR 0.8-1.5), and Graves' disease (RR 2.8). RLN identification (RR 1.6) significantly ($p = 0.01$) reduced permanent RLN palsy rates, unlike parathyroid gland identification during hypoparathyroidism.

Chiang FY et al conducted a study on 107 patients for clinically benign thyroid disease who underwent total thyroidectomies performed by the same surgeon between January 1987 and December 2004 were enrolled; 48 had Graves' disease and 59 had non-Graves' disease.¹² The rates of temporary vs permanent hypoparathyroidism, hematoma requiring surgical intervention, and temporary vs. permanent recurrent laryngeal nerve palsy (RLNP) after total thyroidectomy for benign thyroid disease were 34.6% vs. 3.7%, 6.5%, and 6.5% vs. 1.85%, respectively. The rates of permanent hypoparathyroidism and temporary RLNP in the Graves' disease group were significantly different when compared with the non-Graves' disease group (8.3% vs 0% and 11.5% vs 2.5%, respectively). However, comparing the rates of temporary hypoparathyroidism, permanent RLNP, and postoperative hematoma, there was no statistically significant difference. Compared with total lobectomy, the rates of postoperative hematoma increased significantly for total thyroidectomy (6.5% vs 0.48%).

CONCLUSION

For multinodular goiter disease for excessive bleeding during surgery, a large goiter presented as a predictive factor, and transfusion of blood should be considered in cases in which goiter weighs more than 200 g.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. National Institute of Diabetes and Digestive and Kidney Diseases. Graves' Disease. NIH. 2012. Available at www.niddk.nih.gov.
2. Burch HB, Cooper DS. Management of Graves disease: a review. *JAMA*. 2015;314(23):2544-54.
3. Brent, Gregory A. Clinical practice. Graves' disease. *New Eng J Med*. 2008;358(24):2594-605.
4. Menconi F, Marcocci C, Marinò M. Diagnosis and classification of Graves' disease. *Autoimmunity Rev*. 2014;13(4)(5):398-402.
5. Nikiforov, Yuri E, Biddinger, Paul W, Nikiforova Lester DR, Biddinger, Paul W. Diagnostic pathology and molecular genetics of the thyroid (2nd ed. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2012:69.
6. Yamanouchi K, Minami S, Hayashida N, Sakimura C, Kuroki T, Eguchi S. Predictive factors for intraoperative excessive bleeding in Graves' disease. *Asian J Surg*. 2015;38(1):1-5.
7. Grodski S, Stalberg P, Robinson BG, Delbridge LW. Surgery versus radioiodine therapy as definitive management for Graves' disease: the role of patient preference. *Thyroid*. 2007;17(2):157-60.
8. Erbil Y, Giris, M, Salmaslioglu A. The effect of anti-thyroid drug treatment duration on thyroid gland microvessel density and intraoperative blood loss in patients with Graves' disease. *Surg*. 2008;143:216e225.
9. Karamanakos SN, Markou KB, Panagopoulos K. Complications and risk factors related to the extent of surgery in thyroidectomy. Results from 2,043 procedures. *Hormones*. 2010;9:318e325.
10. Allannic H, Fauchet R, Orgiazzi J. Antithyroid drugs and Graves' disease: a prospective randomized evaluation of the efficacy of treatment duration. *J Clin Endocrinol Metabol*. 1990;70:675e679.
11. Thomusch O, Machens A, Sekulla C. Multivariate analysis of risk factors for postoperative complications in benign goiter surgery: prospective multicenter study in Germany. *World J Surg*. 2000;24:1335e1341.
12. Chiang FY, Lin JC, Wu CW. Morbidity after total thyroidectomy for benign thyroid disease: comparison of Graves' disease and non-Graves' disease. *Kaohsiung J Med Sci*. 2006;22:554e559.

Cite this article as: Sagar RC, Ravikumara G, Malipatil VB. Predictive factors for intraoperative excessive bleeding in multinodular goitre with or without hyperthyroidism. *Int Surg J* 2018;5:3733-7.