Case Report

DOI: http://dx.doi.org/10.18203/2349-2902.isj20194083

Fibroadenoma of ectopic breast tissue in chest wall: a rare case report and review of the literature

Pradeep Balmiki*, Kamlesh Mourya

Department of Surgery, Atal Bihari Vajpayi Government Medical College, Vidisha, Madhya Pradesh, India

Received: 29 May 2019 Revised: 26 July 2019 Accepted: 31 July 2019

*Correspondence:

Dr. Pradeep Balmiki,

E-mail: dr.pradeepkumar2011@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Fibroadenoma of ectopic breast tissue is a rare disease. Ectopuc breast tissues are found along milk line. Malignancy is more common than fibroadenoma in ectopic breast tissue. Ectopic breast tissue can be present with nipple or without nipple areola complex. Here we report a case of 20 years old patient having fibroadenoma of ectopic breast tissue in anterior chest wall without nipple areola complex. Patient underwent excision and biopsy. Histopathology confirmed it as fibroadenoma of the ectopic breast tissue. This case has been reported for its rarity and to reemphasise the importance of screening of EBT for any pathology during routine screening of breast.

Keywords: Ectopic breast tissue, Fibroadenoma, Milk line

INTRODUCTION

Presence of more than two breasts is called as polymastia also often described as supernumerary breast, accessory breast or ectopic breast tissue (EBT). They may be presented either with nipple areolar complex (polythelia) or without nipple areolar complex (polymastia). Ectopic breast tissue can occur anywhere on the primitive embryonic milk line extending from axilla till groin.¹ Axillary breast tissue is a common variant of ectopic breast tissue and is found in approximately two to six percent of general population.² It is twice as common in females when compared to males.3 Carcinoma is more common occurance in ectopic breast tissue followed by inflammatory changes and fibroadenoma.^{3,4} Thus fibroadenoma of ectopic breast tissue is a rare entity and proper evaluation is required to rule out any chances of neoplastic changes.

CASE REPORT

A 20 years old female came to our outpatient department after noticing a swelling below her left breast since 1

month. She was a null gravida. Swelling is not associated with pain, tenderness and local rise of temperature. On Examination, a single 4×3 cm oval shaped swelling was found in subcutaneous plane which is firm in consistency, well demarcated borders and freely mobile in vertical and horizontal planes. Skin over and around the swelling is normal. No discharge or opening found over the swelling. No engorged veins. Bilateral breasts and left axilla is normal. No history of similar swellings or other breast diseases in the family. No bladder and bowel abnormalities. All other systems are normal. Provisional clinical diagnosis of lymphadenopathy was made. Fine needle aspiration cytology was done which suggested the swelling as fibroadenoma of ectopic breast tissue. Routine surgical workup was done which did not reveal any abnormalities. Excision and biopsy was done (Figure 1). Histopathological examination was done in which the sections studied showed mammary ductules surrounded by fibrocollagenous bundles, cystically dilated spaces, foamy macrophages and crowding of gland spaces (adenosis) (Figure 2 and 3). No evidence of malignancy and no lymph node structure identified. Considering the above features and in correlation of the FNAC a diagnosis of "fibroadenoma of ectopic breast tissue" was made.



Figure 1: Specimen is being excised.



Figure 2: FNAC.




Figure 3: Histopathology.

DISCUSSION

Presence of more than two breasts is called as polymastia also often described as supernumerary breast, accessory breast or ectopic breast tissue (EBT). Mammary milk lines represent two ectodermal thickenings which grow on either sides of the embryo during the 6th week of development. They extend from axilla till the groin. As the development precedes most of them disappears except for the two mammary ridges in the pectoral region which later develop into normal breasts. Ectopic breast usually occurs along the "milk line" or mammary line.⁵ Failure of any portion of the mammary ridge to involute can lead to ectopic breast tissue either with nipple areolar complex i.e. polythelia or without nipple areolar complex i.e. polymastia.⁵ Polymastia occurs due to regression of ectodermal element of the skin but a normal proliferation of mesenchyme.6

EBT may occur unilaterally or bilaterally. The incidence of ectopic breast is 1-6% in the general population.⁷ The axillary breast tissue is a subtype of ectopic breast occurring in 2-6% of women.² It is twice as common in female patients as in male patients.³ Approximately 67% of accessory breast tissue occurs in the thoracic or abdominal portions of the milk line often just below the infra mammary crease, more often on the left side of the body and only 20% of accessory breast tissue occur in axilla, the remaining 13% of these occur anywhere along the milk line. Ectopic breast tissues have been reported even on the face, foot, perineum and vulva.8-10 Uncommon locations, also known as 'mammae erraticae', include the buttocks, back of neck, flank, upper arm, hip, shoulders and midline of the back and chest.11

The occurrence of ectopic breast tissues in locations other than the milk line can be explained by two hypothesis. Either it can represent a migratory arrest of breast primordium during chest development or this occurrence can be attributed to its development from the modified apocrine glands.⁶

Carcinomas are common in ectopic breasts, hence it is very important to diagnose the underlying pathology, especially in case of ectopic breasts. Other presentations include cycstic changes, inflammatory changes and fibroadenomas. Ectopic breasts might be associated with an underlying genitourinary and cardiovascular system abnormality. Fibroadenoma of breast is most common cause of breast mass especially in adolscents and young females, however fibroadenoma of ectopic breast tissue is rarely described. Supernumerary breasts are classified by Kajava in 1915 as follows,

- type 1: complete breast with areola and nipple;
- type II: supernumerary breast with nipple only;
- type III: supernumerary breast with breast tissue and areola only;
- type IV: ectopic or aberrant breast tissue only;

- type V: pseudomamma consists of fat with areola and nipple;
- type VI: polythelia (nipples only);
- type VII: areola only (polythelia areolaris)
- and type VIII: patch of hair only (polythelia pilosa). 14

Our case belongs to class IV.

Ectopic breast tissue occurs sporadically, but a hereditary predisposition has also been reported. ¹⁵ In most cases, accessory breasts are asymptomatic and are more of cosmectic concern without any sympotoms or signs. Under circumstances like mensuration, pregnancy and lactation, the size of swelling might increase along with pain which causes discomfort. Carcinoma which commonly occurs from ectopic breast tissue has poorer prognosis due to delay in the diagnosis. This delay happens due to a broad differential diagnosis. Differential diagnosis of a axillary swelling include lipoma, sebaceous cyst, vascular lesions, suppurative hidradenitis, cat scratch disease, lymphadenopathy, secondaries in lymphnodes, tuberculosis, axillary tail of Spence, or even a torn muscle belly and malignancies. 12 Firboadenoma of axillary ectopic breast is present in superficial plane which in contrast to any tumor from axillary tail of Spence which lies deep to the deep fascia. EBT associated with underlying urogenital abnormalities such as hydronephrosis, polycystic kidney and ureteric stenosis. 16,17 This association can be partly explained by the parallel development of mammary structure and genitourinary system.¹⁸ Cardiac problems such as congenital heart anomalies, high-blood pressure and rhythm disturbances are linked with ectopic breasts.

A thorough examination of the swelling has to be done just like in case of any breast disease by triple assessment to rule out any malignancy. Special attention should be given for the examination of lymph nodes as the malignancies of ectopic breast tissues carry poor prognosis. Evaluation also has to be done to rule out any associated urological and cardiac problems. Both normal breasts must be examined. Ultrasound is used to determine the anatomical location, consistency, vascularity and composition of the mass. FNAC or true cut biopsy helps us in arriving at a more accurate provisional diagnosis which helps us in planning of management. If the lesion is found to be benign, a simple excision is sufficient. If the lesion is suspected to be malignant wide excision along with lymph node dissection has to be done following oncological principles.

CONCLUSION

In conclusion, when tumours or nodules are found along the mammary line, the presence of breast tissue should be considered during the investigation. Fibroadenoma of ectopic breast is an extremely rare entity. Proper diagnosis and treatment is essential as the carcinoma of ectopic breast tissue which is common and usually carries a poorer prognosis because of the delayed diagnosis. In our case, excision of the fibroadenoma has been done and the patient is on regular follow up.

ACKNOWLEDGEMENTS

My sincere thanks to our Dean A.B.V.G.M.C Vidisha Dr. Sunil Nandeshwar, Medical Superintendent Dr. Prashant W., HOD Surgery Dr. D. Paramhans, Civil Surgeon Dr. Sanjay Khare for constant encouragement and thanks to Dr. Anup Verma, Dr. P.C. Manjhi and my colleagues Rajkishor Singh, Sandeep Ahirwar and Neeraj Jain for great support. Thanks to Dr. Kamlesh Mourya for constant support, encouragement and help.

Funding: No funding sources Conflict of interest: None declared Ethical approval: Not required

REFERENCES

- 1. Harun C, Ertan B, Ozer A, Rana C. Fibroadenoma of the ectopic breast of the axilla a case report. Pol J Pathol. 2006;57(4):209-11.
- 2. Coras B, Landthaler M, Hofstaedter F, Meisel C, Hohenleutner U. Fibroadenoma of the axilla. Dermatol Surg. 2005;31:1152-4.
- 3. Burdick AE, Thomas KA, Welsh E, Powell J, Elgart GW. Axillary polymastia. J Am Acad Dermatol. 2003;49:1154-6.
- 4. Rizvi G, Pandey H, Gupta MK. Fibroadenoma of ectopic breast tissue in axilla. J Case Reports. 2012;2(2):13–5.
- 5. Shin SJ, Sheikh FS, Allenby PA, Rosen PP. Invasive secretory (juvenile) carcinoma arising in ectopic breast tissue of the axilla: Arch Pathol Lab Med. 2001;125:1372-5.
- 6. Gajaria PK, Maheshwari UM. Fibroadenoma in axillary ectopic breast tissue mimicking lymphadenopathy: J Clin Diagn Res. 2017;11(3):ED01-2.
- 7. Gutermuth J, Audring H, Voit C, Haas N. Primary carcinoma of ectopic axillary breast tissue. J Eur Acad Dermatol Venereol. 2006;20:217–21.
- 8. Koltuksuz U, Aydin E. Supernumerary breast tissue: a case of pseudomamma on the face. J Pediatric Surg. 1997;32(9):1377–8.
- 9. Conde DM, Kashimoto E, Torresan RZ, Alvarenga M. Pseudomamma on the foot: an unusual presentation of supernumerary breast tissue. Dermatol Online J. 2006;12(4):7-10.
- 10. Aughsteen AA, Almasad JK, Al-Muhtaseb MH. Fibroadenoma of the supernumerary breast of the axilla. Saudi Med J. 2000;21:587–9.
- 11. Lilaia C, Capela E, Cadilla JS, Cabrita b. Ectopic breast fibroadenoma. Internet J Gynecol Obstet. 2007;6:2.

- 12. Ciralik H, Bulbuloglu E, Arican O, Citil R. Fibroadenoma of the ectopic breast of the axilla—a case report. Pol J Pathol. 2006;57:209–11.
- 13. Conde DM, Torresan RZ, Kashimoto E, Carvalho LE, Cardoso Filho C. Fibroadenoma in axillary supernumerary breast: case report. Sao Paulo Med J. 2005;123:253–5.
- 14. Kajava Y. The proportions of supernumerary nipples in the Finnish population. Duodecim. 1915;31:143–70.
- 15. Korumilli RK, Srikanth J, Muvva SH, Saka L. Fibroadenoma of ectopic breast tissue in axilla: a case report. Int Surg J. 2018;5:1592-4.
- 16. Varsano IB, Jaber L, Garty BZ, Mukamel MM, Grünebaum M. Urinary tract abnormalities in children with supernumerary nipples. Pediatrics. 1984;73:103–5.
- 17. Mehes K. Association of supernumerary nipples with other anomalies. J Peds. 1979; 95:274.
- 18. Grossl NA. Supernumerary breast tissue: historical perspectives and clinical features. Southern Med J. 2000;93(1):29-32.

Cite this article as: Balmiki P, Mourya K. Fibroadenoma of ectopic breast tissue in chest wall: a rare case report and review of the literature. Int Surg J 2019;6:3389-92.