

## Case Report

# Congenital penile epidermal cyst: a rare location

Abishai Jebaraj\*, Santosh Patil, Vinay S. Kundargi, Basavesh S. Patil

Department of Urology, Shri B. M. Patil Medical College and Research Institute, Bijapur, Karnataka, India

**Received:** 12 January 2020

**Accepted:** 28 January 2020

**\*Correspondence:**

Dr. Abishai Jebaraj,

E-mail: [abishai\\_u@yahoo.co.in](mailto:abishai_u@yahoo.co.in)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

A 22 years old male presented with a mass over the penis since birth, which was painless and gradually increasing in size. The mass was excised and it was found to be penile epidermal cyst. Epidermal cyst is one of the most common benign tumors occurring in the body, which commonly occurs either congenitally or following trauma or surgery, where the epidermal elements get trapped within closed space. But here this case had no trauma and surgeries in the past and the mass present since birth—a case of congenital penile epidermal cyst.

**Keywords:** Penile cyst, Epidermal cyst, Congenital, Rare presentation

### INTRODUCTION

Epidermal cyst can occur in any part of the body.<sup>1</sup> It occurs due to implantation of epidermal elements in the dermis during the fetal formation, trauma or surgeries like penoplasty, circumcision, etc. It can also result from the proliferation of epidermal cells within a circumscribed space of the dermis.<sup>2</sup> Penile epidermoid cysts are not very common and usually congenital.<sup>3</sup> The etiology in penile area is not specific, but it may be due to an abnormal embryogenic closure of the median raphe or a monolayer teratoma of germ cell origin.<sup>4</sup> We report a case of a congenital penile cyst in a 22 years old male without any history of penile surgery and injury.


### CASE REPORT

A 22 years old male presented to the outpatient department of urology, with a swelling over the penis since birth. It was painless and was gradually increasing in size. There was no past history of surgery/local trauma/infection. There were no associated urinary complaints.

On local examination, single mass measuring 3×2 centimeters, firm in consistency, globular in shape and non-tender on palpation was noted in the ventral aspect of the penis in the subcutaneous plane, arising from the penile shaft skin. There were no visible scars or punctum noted over it (Figure 1).



**Figure 1:** Epidermal cyst of penis (preoperative image).



**Figure 2: Ultrasound image showing hypoechoic lesion with hyperchoic foci.**

Ultrasonography of the penile mass reported, a 3.2×1.6 centimeters sized well defined predominantly hypoechoic lesion with multiple hyperchoic foci showing comet tail artifact, noted in subcutaneous plane on ventral side of penis. There was evidence of vascularity in the periphery of the lesion likely arising from the frenulum. However, there was no vascularity inside the lesion. The features were suggestive of epidermal cyst of penis (Figure 2).




**Figure 3: Excised specimen of congenital penile epidermal cyst.**

Patient underwent wide local excision of the lesion. Swelling was not adherent to the underlying structures like urethra or frenulum and the post-operative period was uneventful (Figure 3).

Histopathology of the excised specimen revealed tissue lined by stratified squamous epithelium. Sub epithelium showed cyst wall lined by stratified squamous epithelium with prominent granular layer with the lumen of the cyst

showing lamellated keratin suggestive of epidermal cyst (Figure 4).



**Figure 4: Histopathology showing (a) lamellated keratin and (b) prominent granular layer.**

## DISCUSSION

Epidermal cysts are benign lesions occurring in any part of the body. They are more common in men.<sup>5</sup> Epidermal cyst of penis is a rare entity and is either congenital or acquired. Amarnath et al in their study have reported that there were less than 200 cases of epidermal cyst of penis were reported in the past of which only ten cases were reported from India and among that only one case of congenital penile epidermal cyst was reported.<sup>6</sup>

Various etiologies have been proposed for the formation of the cyst which includes sequestration of epidermal rests during embryonic life, traumatic or surgical implantation of epithelial elements like after circumcision.<sup>7</sup>

They are usually asymptomatic during childhood but become symptomatic during adulthood. The usual symptoms of penile epidermal cyst are local pain, difficulty in micturition, haematuria, haemospermia and difficulty in having sexual intercourse. The differential diagnosis of eridermoid cyst of penis includes dermoid cyst, teratoma, glomustumour, pilonidal cyst and steatocystoma.<sup>7</sup>

Infection, dyspareunia, urine flow obstruction and cosmesis are some of the indications of treatment of epidermal cyst.<sup>3</sup> Though malignancies arising from epidermal cysts in other parts of body are reported, there are no such reports from penile epidermal cysts so far in the literature.<sup>8</sup>

## CONCLUSION

Our patient most probably had developed the congenital epidermal cyst during his embryonic life. Simple excision

and follow-up are the line of management as the risk of malignancy is very low. Reports of congenital penile cysts are very few in literature. Only one case of congenital penile epidermal cyst has been reported in India, till now and our case report will add up to the list of such rare case.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: Not required*

## REFERENCES

1. Singh S, Kaur T. Epidermoid cyst of penis. Indian J Dermatol Venereol Leprol. 2011;77:627.
2. Saini P, Mansoor MN, Jalali S, Sharma A. Penile epidermal inclusion cyst. Indian J Pediatr. 2010;77:815-6.
3. Suwa M, Takeda M, Bilim V, Takahashi K. Epidermoid cyst of the penis: A case report and review of the literature. Int J Urol. 2000;7:431-3.
4. Okumura A, Takagawa K, Fuse H. Penile epidermoid cyst consisted of multiple foci: A case report. J Cosmet Dermatol Sci Appl. 2012;2:214-5.
5. Hanson L, Zeitouni N. Epidermal inclusion cyst. EMedicine Clin Knowledge Base. 2009;1061582.
6. Amaranathan A, Sinhasan SP, Dasiah SD. Median raphe cysts of the prepuce skin, with triple histological linings: a case report and review of the literature. J Clin Diagn Res. 2013;7(7):1466-8.
7. Chandanwale SS, Buch AC, Kumar H, Mishra N. Epidermoid cyst in the breast: A common benign lesion at a rare site. Clin Cancer Investig J. 2015;4:99-101.
8. Romani J, Barnadas MA, Miralles J, Curell R, de Moragas JM. Median raphe cyst of the penis with ciliated cells. J Cutan Pathol. 1995;22:378-81.

**Cite this article as:** Jebaraj A, Patil S, Kundargi VS, Patil BS. Congenital penile epidermal cyst: a rare location. Int Surg J 2020;7:889-91.