

## Original Research Article

# Study of the association of serum procalcitonin with leukocytosis in predicting complicated acute appendicitis

Thana Ram Patel\*, Amit Jain, Lakshman Agarwal, Vakta Ram Choudhary, Dinesh Bijarniya

Department of General Surgery, SMS Medical College, Jaipur, Rajasthan, India

**Received:** 06 May 2020

**Revised:** 11 June 2020

**Accepted:** 12 June 2020

**\*Correspondence:**

Dr. Thana Ram Patel,

E-mail: [dr.thanarampatel@gmail.com](mailto:dr.thanarampatel@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** Delay in the diagnosis of complicated appendicitis and its treatment results in an increased rate of postoperative morbidity, mortality and hospital stay. The diagnosis of appendicitis rests on a combination of signs of inflammation such as fever, pain and tenderness; leukocytosis, and elevated C-reactive protein levels, interleukin-6 (IL6) and procalcitonin. Raised level of serum procalcitonin in bacterial infection has been used to further improve the diagnosis of complicated AA.

**Methods:** One-hundred ten patients of appendicitis confirmed by intra-operative findings and final pathologist report, who underwent appendectomy consisting 25 women (22.73%) and 85 men (77.27%) with a mean age of 25 years (age range 15-55 years) were included in this study. Serum procalcitonin value was measured by chemi E411 Cobas method (chemiluminescent immunoassay system) using the B.R.A.H.M.S PCT kit. Serum PCT level >0.5 ng/ml was consider as risk for progression to severe systemic disease.

**Results:** At a 0.5 ng/dl cut-off value of PCT, the sensitivity and specificity of PCT level measurement for acute complicated appendicitis prediction was 90% and 97.14% respectively. Association between WBC count and PCT value shows the sensitivity and specificity in 40 case of acute complicated appendicitis prediction was 86% and 75% respectively.

**Conclusions:** Both the higher PCT values and leukocytosis correlates well with infectious post-operative complications for acute appendicitis and it can help to carry out timely surgical intervention which is highly recommended in complicated appendicitis (correlates PCT >0.5 ng/dl).

**Keywords:** Complicated AA, Leukocytosis, Procalcitonin

### INTRODUCTION

It is estimated that as much as 6% to 7% of the general population will develop appendicitis during their lifetime, with the incidence peaking in the second decade of life.<sup>1</sup>

As appendicitis progresses, the blood supply is impaired by bacterial infection in the wall and distention of the lumen; gangrene and perforation occurs at about 24 hours, though the timing is highly variable. Gangrene implies microscopic perforation, bacterial contamination

of the peritoneum, and peritonitis. This process may be effectively localized by adhesions from nearby viscera.<sup>2</sup>

Appendectomy remains one of the most commonly performed operations worldwide.<sup>3</sup>

The clinical diagnosis of appendicitis rests on a combination of localized pain and tenderness accompanied by signs of inflammation, such as fever, leukocytosis, and elevated C-reactive protein levels. In the absence of signs of inflammation the diagnosis is less certain, and in this situation a CT scan may be of value.<sup>2</sup>

Following the introduction and widespread use of antibiotics in the 1940s, mortality rates improved further. C-reactive protein, bilirubin, Il-6, and procalcitonin have all been suggested to be helpful in the diagnosis of appendicitis, specifically in predicting perforated appendicitis.<sup>4</sup>

Delay in diagnosis and treatment results in an increased rate of perforation, postoperative morbidity, mortality and hospital stay.

**Aim**

The aim of this study was to assess the value of procalcitonin as a predictor of severity of appendicitis in order to improve the clinical decision making.<sup>5,6</sup>

**Objectives**

*Primary objective:* To assess the value of procalcitonin as a predictor of severity (inflamed, gangrenous, perforated) of appendicitis in order to improve the clinical decision making for timely intervention.

*Secondary objective:* To establish relation between PCT, leukocytosis histopathological finding of appendix and its correlation with post-operative complications.

**METHODS**

Study design was hospital based cross sectional study. Study was conducted in between 1 January 2018 to 31 November 2019. Study area was Department of general surgery of SMS hospital, SMS medical college, Jaipur.

Study population was selected as per the inclusion and exclusion criteria.

**Inclusion criteria**

Those patients who have given written and informed consent, Age group 15-50 years in either sex, all patients without exclusion criteria, who underwent appendectomy, in which peri-operatively complicated appendicitis were found.

**Exclusion criteria**

Diagnosis other than acute appendicitis confirmed during the operation, Presence of another acute infectious lesion or disease in any other part of the body, immunosuppressive disease like HIV.

Sample size was calculated at 95% confidence level assuming 65% specificity of PCT to detect peritonitis due to appendicitis and prevalence of peritonitis due to appendicitis was 9.5%. At 10% absolute allowable error sample size was found 102 patients of peri-operatively complicated confirmed appendicitis which was further rounded off to 110 such patients.

**Methodology**

Patients diagnosed with symptomatic appendicitis were selected based on the clinical signs or USG report. The patients were informed about the details of the procedure and written informed consent was obtained. All routine blood tests specifically TLC and serum procalcitonin were assessed. And all the cases divided in two groups on the basis of procalcitonin value. Serum procalcitonin value was measured by chemi E411 Cobas method (chemiluminescent immunoassay system) using the B.R.A.H.M.S PCT kit.

Status of the appendix (peri-operative finding) and histological diagnosis (normal/ inflamed/ gangrenous/ perforated) was recorded. Patients were analyzed whether they have uncomplicated or complicated appendicitis, complicated appendicitis patients were those having gangrenous or perforated appendix with or without appendicular abscess formation. Serum procalcitonin value <0.5 ng/ml was considered normal. Patients with procalcitonin value >0.5 ng/ml (value as a risk for progression to severe systemic disease) were consider positive. Similarly TLC >10,000 was taken significant.

Statistical analysis was performed by MS excel worksheet 2007. Analysis of distribution of data was assessed by the Chi-square test.

**RESULTS**

One-hundred ten patients of appendicitis confirmed by intra-operative findings and final pathologist report, who underwent appendectomy consisted of 25 women (22.73%) and 85 men (77.27%) with a mean age of 25 years (age range 15-55 years) were included in this study.


Mean age was 25 years with standard deviation 31.8. Age group 15-24 years and 25-34 years had the most common incidence of appendicitis.

**Table 1: Age and sex wise distributions of the acute appendicitis case.**


Age/sex	Female	Male	Total
	N (%)	N (%)	N (%)
15-24	15 (60)	53 (62.35)	68 (61.82)
25-34	6 (24)	19 (22.35)	25 (22.73)
35-44	1 (4)	8 (9.41)	9 (8.18)
45-55	3 (12)	5 (5.88)	8 (7.27)
15-55	25 (100)	85 (100)	110

Female patients having uncomplicated appendicitis were 18 (72%) and complicated in 7 (28%) patients while male patients having uncomplicated appendicitis were 52 (61.18%) as compared to complicated appendicitis in 33 (38.82%) patients.

Procalcitonin value in 73 patients (26.03% female and 73.97% male patients) was less than 0.5 ng/ml and thence considered negative. In the remaining 37 patients (16.22% female and 83.78% male patients), serum PCT value was more than 0.5 ng/ml and these considered positive (Table 2).



**Figure 1: Sex wise distribution of complicated and uncomplicated acute appendicitis.**



**Figure 2: Distribution of complicated and uncomplicated acute appendicitis.**

**Table 2: Sex wise distribution of cutoff level of procalcitonin in acute appendicitis**

Gender/PCT	Uncomplicated (AA<0.5)	Complicated (AA>0.5)
	N (%)	N (%)
Female	19 (26.03)	6 (16.22)
Male	54 (73.97)	31 (83.78)
Total	73 (66.36)	37 (33.64)

Based on intraoperative finding with the final pathologic report, we classified the patients into two groups: Group I

with a complicated appendicitis and Group II with uncomplicated appendicitis.

At a 0.5 ng/dl cut-off value of PCT, the sensitivity and specificity of PCT level measurement for acute complicated appendicitis prediction was 90% and 97.14% respectively. The PPV and NPV of PCT in complicated appendicitis were (94.74%) and (94.44%) respectively.

**Table 3: Association of PCT with complicated AA.**

PCT/AA	Complicated AA	Uncomplicated AA	Total
>0.5	36	2	38
<0.5	4	68	72
	40	70	110

Chi-square =81.676 with 1 degree of freedom; P<0.001  
LS=Significant

The severity of the appendix inflammation determined for each patient at the time of operation and confirmed by histopathological report known as complicated appendicitis included the following: abscess/gangrenous/perforated appendix (40 patients). Complicated appendicitis may have abscess or perforation (peritonitis) or necrotizing appendicitis which may lead to frequent post-operative complication like surgical site infection.

Out of 18 patients with necrotizing appendicitis, 16 (88%) patients had PCT level more than 0.5 ng/ml while only 22 (23%) patients out of 92 were having high levels of PCT in uncomplicated appendicitis group.

Peritonitis due to perforated appendicitis was confirmed in 25 patients intra-operatively. Among 25 patients with peritonitis, 21 patients had a PCT value more than 0.5 ng/ml.

Post-operative complications including surgical site infection (SSI) was detected in 8 patients. In 8 cases of surgical site infection, 7 patient had a PCT value more than 0.5 ng/ml, and one patient had a negative PCT value.

The corresponding sensitivity for detecting peritonitis, necrotizing appendicitis and surgical site infection were 84%, 88.89% and 87.5% and their specificity were 80%, 76.09% and 69.61% respectively.

**Table 4: Association of WBC and PCT in 40 acute complicated appendicitis cases.**

WBC\PCT	>0.5	<0.5	Total
>10,000	31	1	32
<10,000	5	3	8
	36	4	40

Chi-square=8.4 with 1 degree of freedom; P<0.05

Association between WBC count and PCT value shows the sensitivity and specificity in 40 case of acute complicated appendicitis prediction was 86% and 75% respectively.

All 40 cases had Alvarado score >7 is strongly suggestive of acute appendicitis, using definition leukocytosis (>10,000 WBC/ $\mu$ l).<sup>7</sup>

## DISCUSSION

This study included 110 patients with 25 females and 85 males with a mean age of 25 years.

Based on the intra-operative findings (abscess, gangrenous or perforated appendix), each case was labelled as uncomplicated appendicitis or complicated appendicitis. Peritonitis due to perforated appendicitis was confirmed in 25 patients intra-operatively.

Post-operative complications including surgical site infection (SSI) was detected in 8 patients.

Procalcitonin value in 72 patients was less than 0.5 ng/ml and hence considered negative. In the remaining 38 patients, serum PCT value more than 0.5 ng/ml.

Among 25 patients with peritonitis, 21 patients had a PCT value more than 0.5 ng/ml. In 8 cases of surgical site infection, 7 patient had a PCT value more than 0.5 ng/ml, and one patient had a negative PCT value.

At a 0.5 ng/dl cut-off value of PCT, the sensitivity and specificity of PCT level measurement for acute complicated appendicitis prediction was 90% and 97.14% respectively.

The PPV and NPV of PCT in complicated appendicitis were (94.74%) and (94.44%) respectively

The corresponding sensitivity for detecting peritonitis, necrotizing and surgical site infection were 84%, 88.89% and 87.5% and their specificity were 80%, 76.09% and 69.61% respectively.

Chandel et al in their study of 40 patients found sensitivity (95.65%), specificity (100%) and predictive values PPV (100%) and NPV (83.3%) of serum PCT levels as the diagnostic test for the diagnosis of acute appendicitis.<sup>8</sup>

Gavela et al study done for diagnosis of peritonitis, a PCT cutoff of 0.18 ng/ml gave a sensitivity of 97%, specificity of 80%, positive predictive value of 72%, and negative predictive value of 89.3%.<sup>9</sup>

In Kafetzis et al study, there was a statistically significant indication ( $p < 0.0001$ ) that PCT levels >0.5 ng/ml were observed more frequently in cases of perforated or necrotizing appendicitis than in simple appendicitis.<sup>10</sup> A

PCT level of >0.5 ng/ml can indicate perforation or gangrene with 73% sensitivity and 94% specificity.

In Kouame et al study of 101 children, the sensitivity of PCT was 28%: 95% CI (18-40), 88% specificity: 95% CI (72-97), the predictive value 83% positive 95% CI (61-95) and predictive value negative 37%: 95% CI (26-49).<sup>11</sup> The sensitivity does not allow a threshold value to be determined for diseased appendages. There is a positive correlation between the serum PCT level greater than 0.5  $\mu$ g/l and severe appendicular lesions. PCT mean increases with the severity of inflammatory lesions of the appendix ( $p = 0.0051$ ).

Sand et al noted that the sensitivity of PCT for the diagnosis of acute appendicitis was 0.14, with a specificity of 1.<sup>12</sup> The PPV of PCT (1-0.5 ng/ml) for appendicitis was 1 in contrast to 0.76 for CRP (1-0.5 mg/dl) and to 1 for WBC. The NPV for appendicitis were 0.06 for PCT, 0.09 for WBC and 0.11 for CRP. The values of CRP, WBC and PCT increased with the severity of the appendicitis.

In Wu et al study, the mean age of the study sample was  $35.2 \pm 18.3$  years old and 110 patients (51.4%) were male.<sup>13</sup> In diagnosing appendicitis and complicated appendicitis the AUC were 0.69 and 0.76 for PCT respectively.

Yamashita et al consider five parameters- plasma procalcitonin level (PCT), body temperature (BT), white blood cell count (WBC), neutrophil/lymphocyte ratio (N/L ratio), and C-reactive protein level (CRP).<sup>14</sup> Univariate analysis of the predictors of abscess and/or perforation revealed that a plasma PCT level  $\geq 0.46$  ng/mL had the highest odds ratio (30.3 [95% confidence interval: 6.5-140.5] versus PCT <0.46 ng/ml). These findings indicate that procalcitonin is a useful marker of acute appendicitis with abscess and/or perforation.

Yu et al study ROC curve analysis showed that WBC had high accuracy (area under ROC curve 0.72 per cent C.I. 0.68 to 0.76), followed by procalcitonin (0.65, 0.61 to 0.69).<sup>15</sup> Procalcitonin was found to be more accurate in diagnosing complicated appendicitis, with a pooled sensitivity of 62 (33 to 84) per cent and specificity of 94 (90 to 96) per cent.

In present study WBC count and PCT value shows the sensitivity and specificity in 40 case of acute complicated appendicitis prediction was 86% and 75% respectively.

## CONCLUSION

In complicated acute appendicitis leukocytosis suggest further PCT measurement. The PCT values can be used as predictor of complicated appendicitis and infectious complications following surgery for acute appendicitis and it can help to carry out timely surgical intervention which highly is recommended in patients with values

more than 0.5 ng/ml and increased PCT values correlates well with leukocytosis and also in this high litigation era both values can guide you to explain the incidence of possible surgical site complication in complicated appendicitis to the patients and its attendants.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

1. Richmond B. The Appendix. In: Townsend Jr M, Beauchamp RD, Evers BM, Mattox KL, eds. Sabiston textbook of surgery: the biological basis of modern surgical practice. 20th edn. Philadelphia: Elsevier; 2017:1296
2. Pennington EC, Burke PA. Appendix. In: Gerard M. Doherty GM, ed. Current diagnosis and treatment surgery. 14th edn. New York: McGraw-Hill Education; 2015:651.
3. Velanovich V. The Appendix. In: Josef E. Fischer JE, ed. Fischer's mastery of surgery. 7th edn. Philadelphia: Wolters Kluwer; 2019:4916.
4. Dahdaleh FS, Heidt D, Turaga KK. The appendix. In: Brunnicardi FC, ed. Schwartz's principles of surgery. 11th edn. New York: McGraw-Hill Education; 2019:1331.
5. Meisner M. Introduction. In: Procalcitonin a new, innovative infection parameter biochemical and clinical aspects. 3rd revised and extended edition. New York: Thieme; 2000:1.
6. Assarsson J, Körner U, Lundholm K. Evaluation of procalcitonin as a marker to predict antibiotic response in adult patients with acute appendicitis: a prospective observational study. Surg Infect. 2014;15(5):601-5.
7. Alvarado A. A practical score for the early diagnosis of acute appendicitis. Ann Emerg Med. 1986;15(5):557-64.
8. Chandel V, Batt SH, Bhat MY, Kawoosa NU, Yousuf A, Zargar BR. procalcitonin as the biomarker of inflammation in diagnosis of appendicitis in pediatric patients and prevention of unnecessary appendectomies. Indian J Surg. 2011;73(2):136-41.
9. Gavela T, Cabeza B, Serrano A, Casado-Flores J. C-reactive protein and procalcitonin are predictors of the severity of acute appendicitis in children. Pediatr Emerg Care. 2012;28(5):416-9.
10. Kafetzis D, Velissariou I, Nikolaides P, Sklavos M, Maktabi M, Spyridis G, et al. Procalcitonin as a predictor of severe appendicitis in children. Eur J Clin Microbiol Infect Dis. 2005;24(7):484-7.
11. Kouame DB, Garrigue MA, Lardy H, Machet MC, Giraudeau B, Robert M. Is procalcitonin able to help in pediatric appendicitis diagnosis? Ann Chir. 2005;130:169-74.
12. Sand M, Trullen XV, Bechara FG, Pala XF, Sand D, Landgrafe G, et al. A prospective bicenter study investigating the diagnostic value of procalcitonin in patients with acute appendicitis. Eur Surg Res. 2009;43(3):291-7.
13. Wu JY, Chen HC, Lee SH, Chan RC, Lee CC, Chang SS. Diagnostic role of procalcitonin in patients with suspected appendicitis. World J Surg. 2012;36(8):1744-9.
14. Yamashita H, Yuasa N, Takeuchi E, Goto Y, Miyake H, Miyata K, et al. Diagnostic value of procalcitonin for acute complicated appendicitis. Nagoya J Med Sci. 2016;78(1):79.
15. Yu CW, Juan LI, Wu MH, Shen CJ, Wu JY, Lee CC. Systematic review and meta-analysis of the diagnostic accuracy of procalcitonin, C-reactive protein and white blood cell count for suspected acute appendicitis. Br J Surg. 2013;100(3):322-9.

**Cite this article as:** Patel TR, Jain A, Agarwal L, Choudhary VR, Bijarniya D. Study of the association of serum procalcitonin with leukocytosis in predicting complicated acute appendicitis. Int Surg J 2020;7:2300-4.