Original Research Article

DOI: https://dx.doi.org/10.18203/2349-2902.isj20213119

Evaluation of post-operative complications associated with tobacco use in thoracic surgery patients

Mehmet Degirmenci¹, Celal Kus²*

¹Department of Thoracic Surgery, Kahramanmaras Sutcu Imam University, School of Medicine, Kahramanmaras, Turkey

Received: 17 May 2021 Revised: 04 July 2021 Accepted: 05 July 2021

*Correspondence: Dr. Celal Kus.

E-mail: ckus46@hotmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Tobacco can make thoracic diseases more complicated by affecting their respiratory functions. Smoking causes many diseases that require surgical treatment and affects surgical results. The aim of the study was to determine the relationship between tobacco use and post-operative complications in thoracic surgery patients and contribute to public health.

Methods: In this study, 754 patients were evaluated retrospectively. Patient characteristics and tobacco use habits of the patients were determined. Postoperative complications, admission to the intensive therapy unit, intubation, death, and length of stay in hospital were defined as surgical outcomes. These results were compared and analyzed with tobacco use.

Results: The patients consisted of 536 (71.1%) men and 218 (28.9%) women. Tobacco use was more common in men (X^2 =223.216, p<0.001) and younger ages (X^2 =45.342, p<0.001). Complications occurred in 96 patients, 76 (79.2%) of whom used tobacco. Tobacco use (p<0.001, OR=3.547), ASA score (p=0.029, OR=2.004), major surgeries (p<0.001, OR=4.458), and minimally invasive surgeries (p=0.027, OR=2.323) are associated with complications. Length of hospital stay is related to the amount of tobacco (p<0.001, OR=3.706), size of surgery (p<0.001, OR=14.797), over 65 years (p<0.001, OR=2.635), and infectious diseases (p=0.039, OR=1.939).

Conclusions: Tobacco use is related to poor outcomes in thoracic surgery patients, and it is a severe health problem, especially at young ages. Tobacco control programs should be supported to prevent the effects of tobacco use on thoracic diseases and postoperative complications.

Keywords: Tobacco use, Post-operative complications, Thoracic surgery, Tobacco products, Smoking

INTRODUCTION

Tobacco use is one of the causes of preventable mortality and morbidity. Due to tobacco use in the world, 5.4 million people die every year. Smoking is a public health problem, and 20% of deaths in developed countries are associated with smoking. In Turkey, tobacco use affects about a quarter of deaths. People in the study area use a variety of tobacco products. These include cigarettes,

wrapping tobacco, electronic cigarettes, hookahs, chewing tobacco, and sublingual tobacco powder.⁴

The supply of oxygen, which is indispensable for our body to work, is done by the respiratory system, and the respiratory system is one of our most vital systems.⁵. While smoking causes many diseases that require surgical treatment, especially thoracic surgery, on the other hand, it affects surgical results.⁶ In addition to chronic changes

²Department of Family Medicine, Kahramanmaras Sutcu Imam University, School of Medicine, Kahramanmaras, Turkey

in thoracic surgery patients, acute effects also occur due to the intervention. Therefore, pulmonary complications are more complicated in thoracic surgery patients than in other patients.

The aim of the study was to determine the relationship between tobacco use and postoperative complications and prognosis in thoracic surgery patients. Thus, it was aimed to protect public health by contributing to the prevention of mortality and morbidity related to smoking.

METHODS

This was a retrospective study. This study was conducted in Adiyaman Training and Research Hospital, Kahramanmaras Necip Fazil City Hospital, and Kahramanmaras Sutcu Imam University Health Research and Application Hospital from October 2006 to October 2019.

Inclusion criteria

We scanned files of 2169 patients treated inpatients at thoracic surgery clinics in three different centers. Of the patients, 754 between the ages of 18-95 years were included in the study.

Exclusion criteria

Diseases due to external factors such as blunt and penetrating chest trauma were excluded from the study. Patients under the age of 18 were not included in the study.

Procedure

Information about the tobacco use habits of the patients was obtained from the records. The tobacco products used by the patients and the daily amount of tobacco were learned from the records. The tobacco products used by the patients were cigarettes, electronic cigarettes, wrapped tobacco, hookahs, tobacco chewing, and sublingual tobacco powder. Patients who have not used tobacco products for more than one year are considered former users. The daily number of cigarettes used for cigarette users and electronic cigarette users was calculated. Since other tobacco products do not have standard measurements, they are classified as 'light users' and 'heavy users'. For cigarette users and electronic cigarette users, less than 20 cigarettes per day were considered 'light users', and 20 and above 'heavy users'.

Since the number of patients using tobacco products other than cigarettes was low, tobacco products were classified as 'cigarettes' and 'other tobacco products' while performing the analysis.

Demographic data of the patients were evaluated. The patients were classified according to their gender. The patients were divided into three groups according to their age: 34 years and under', '35-64 years' and '65 years and

over. Treatment methods were divided into three groups as minor surgeries, minimally invasive surgeries, and major surgeries. Patients were divided into diagnostic groups. Comorbidities were determined. The physical performance status of the patients was classified according to the American Society of Anesthesiologists (ASA) score.⁷

Patients were analyzed according to the development of complications. Patients with complications were divided into tobacco users and non-tobacco users. The patients were analyzed according to the tobacco products they used and the amount of tobacco.

We did not include the intensive therapy unit (ITU) admissions planned for postoperative periods. Emergency admission to the intensive care unit, reintubation, and death were considered poor prognosis criteria. Age, gender, tobacco use, diagnostic groups, types of surgery, and ASA status were considered as possible variables associated with these results. These variables were evaluated by logistic regression analysis.

The mean value of the length of stay in hospital was calculated. The patients were divided into two groups according to their hospital stay. Hospital stays shorter than the mean value was considered as short stays, and those staying longer than the mean value as long stays. Patients in each hospitalization group were evaluated according to their tobacco use status. To more accurately evaluate the relationship between tobacco use and hospital stay, logistic regression analysis was performed with other independent variables such as gender, age groups, types of surgery, ASA status, diagnostic groups, and comorbidities.

Ethical approval

The Clinical Research Ethics Committee of Kahramanmaras Sutcu Imam University Faculty of Medicine approved this study with the decision 02 on 18 March 2020.

Statistical analysis

Statistical analysis was performed using SPSS 20.0 for Windows (SPSS, Inc.; Chicago, USA) package program. In the study, mean±standard deviation, median, and minimum-maximum values for continuous numerical variables and number (N) and percentage (%) for categorical variables were used to present descriptive statistics. The Chi square test or Fisher's exact test was used to compare categorical variables in group comparison. The conformity of the obtained data to normal distribution was examined by Kolmogorov-Smirnov/Shapiro-Wilk tests. Normally distributed continuous variables were compared with parametric tests (paired-sample t-test and independent groups t-test), and non-normally distributed continuous variables were compared with nonparametric tests (Mann-Whitney U and Kruskal Wallis test). Post-hoc tests were performed when necessary. The effects of variables (p<0.25) determined by pairwise comparisons in predicting the presence of poor prognosis criteria and length of hospital stay were evaluated by logistic regression analysis. 'The enter method' was used in the logistic regression analysis. Hosmer-Lemeshow test was used to evaluate model fit. The statistical significance limit value was considered as a p value less than 0.05 (p<0.05), and the confidence interval was 95%.

RESULTS

The 754 patients included in the study consisted of 423 (56.1%) tobacco users and, 331 (43.9%) tobacco nonusers. Fifty-one patients (12.05%) of the tobacco users were former tobacco users, while 131 (30.97%) of the current users were light users and 241 (56.98%) were heavy-tobacco users. As tobacco products, 390 (92.2%) of the patients were using cigarettes, 17 (4.0%) of the patients were using sublingual tobacco powder, 9 (2.1%) of the patients were using wrapped tobacco, 4 (1.0%) of the patients were using hookahs, and 3 (0.7%) of the patients were using electronic cigarettes. Thus, 33 (7.8%) patients used tobacco products other than cigarettes. The patients in the study consisted of 536 (71.1%) males and 218 (28.9%) females. Table 1 shows the patient characteristics

of tobacco use. While the rate of tobacco use in males was 73.3%, it was 13.8% in females. This data showed that tobacco use was significantly more common in the male gender (X^2 =223.216, p=0.001) (Table 1). The mean age of patients was 46.91±22.07 (min: 18, max: 95). According to this study, tobacco use was more common in the 34 and under age group (X^2 =45.342, p=0.001) (Table 1).

Comorbidity was present in 198 (26.25%) of the patients (Table 1). The most common chronic disease was congestive heart failure in 83 patients. Other common comorbidities were hypertension in 49 patients, chronic obstructive pulmonary disease in 47 patients, diabetes mellitus in 43 patients, and chronic renal failure in 24 patients. Of the patients with comorbidities, 23 (11.6%) were former users, 22 (11.1%) were light users, 35 (17.7%) were heavy users, and 118 (59.6%) were not using tobacco. There was no significant relationship between tobacco use and comorbidities. There were 202 patients in the ASA I group, 456 patients in the ASA II group, 60 patients in the ASA III group, and 36 patients in the ASA IV group. The distribution of tobacco use, according to the ASA status of the patients, is shown in Table 1. There was no significant relationship between tobacco use and ASA status.

Table 1: Patient characteristics of tobacco use.

Characteristics		Tobacco	Tobacco amounts						
		Non- users	Former users	Light users	Heavy users	Total	\mathbf{X}^{2^*}	P* value	
	Male	N	143	47	114	232	536	226.759	0, 001
Gender		%	26.7	8.7	21.3	43.3	100,0		
Gender	Female	N	188	4	17	9	218		
	remaie	%	86.2	1.9	7.8	4.1	100, 0		
	-21	N	97	6	68	129	300	_	0, 001
	≤34	%	32.3	2.0	22.7	43.0	100, 0	45.342	
	35-64	N	109	18	48	80	255		
Age (years)		%	42.7	7.1	18.8	31.4	100.0		
	≥65	N	125	27	15	32	199		
		%	62.8	13,6	11.5	16.1	100.0		
	1-2	N	291	41	119	207	658	4.210	0.240
A C A & -4-4		%	44.2	6.2	18.1	31,5	100, 0		
ASA† status	≥3	N	40	10	12	34	96		
		%	41.7	10.4	12.5	35.4	100.0		
	Minor	N	230	39	89	157	515		0.525
		%	44.7	7.5	17.3	30,5	100, 0		
Surgery	Minimally	N	31	7	17	23	78	0 (17	
type	invasive	%	39, 7	9.0	21.8	29.5	100.0	0, 617	0, 735
	Major	N	70	5	25	61	161	_	
		%	43.3	3,0	15.2	37.8	100.0		
Tatal	m 1		331	51	131	241	754		
Total		%	43.9	6.7	17, 4	32.0	100.0		

Note: *Calculated by the Chi-square test or the Fisher exact test. †American Society of Anesthesiologist.

We classified the patients according to diagnosis. Two hundred and seventy-four (36.3) patients were in the spontaneous pneumothorax group, 264 (35.0%) patients

were in the benign diseases, 125 (16.6%) patients were in the malign diseases, and 91 (12.1%) patients were in the infectious diseases group. The tobacco use rates of the

patients in each surgical treatment group were compared, and no relationship was found between tobacco use and the type of surgery. Complications developed in 96 of 754 patients included in the study (12.7%). The number of tobacco users in patients with complications is significantly higher than in non-tobacco users ($X^2=23.765$, p<0.001). Tobacco use rates were significantly higher in patients who developed persistent air leaks, atelectasis, and pneumothorax as complications (Table 2). Of the 96 patients who developed complications, 20 (8.3%) did not use tobacco products, 72 (75.0%) were using cigarettes, and 4 (4.2%) were using other tobacco products.

There was a significant difference between the groups $(X^2=24.865, p<0.001)$. When the groups were compared in pairs, it was observed that the difference was due to the difference between cigarette users and non-tobacco users. There was no difference between cigarettes and other tobacco products in terms of complication development.

Of the 96 patients with complications, 8 (21.6%) were former users, 17 (17.7%) were light- users, 51 (53.1%) were heavy-users, and 20 (20.8%) were non-tobacco user.

Table 2: Relationship of complications with tobacco use.

Complications	Tobacco+		Tobacco-		Total		X2*	D* l
Complications	N	%	N	%	N	%	Λ^{-}	P* value
Persistent air leak	23	79.3	6	20.7	29	100.0	6.597	0.010
Atelectasis	21	87.5	3	12.5	24	100.0	9.924	0.002
Pneumothorax	16	80.0	4	20.0	20	100.0	4.765	0.038
Respiratory failure	3	60.0	2	40.0	5	100.0	0,031	0.998
Pleural effusion	5	100.0	0	0.0	5	100.0	3.939	0.071
Surgical site infection	3	75.0	1	25.0	4	100.0	0.583	0.635
Cardiac arrhythmia	2	50.0	2	50.0	4	100.0	0.061	0.999
Pneumonia	1	33.3	2	66.7	3	100.0	0.634	0.585
Acute renal failure	1	100.0	0	0.0	1	100.0	0.784	0.998
Bronchospasm	1	100.0	0	0.0	1	100.0	0.784	0.998
Total	76	79.2	20	20.8	96	100.0		

Note: *Calculated by the Chi-square test or the Fisher exact test.

Table 3 includes the evaluation of relationships of independent variables with the development of complications using logistic regression analysis. Compared to non-smokers, the risk of complications was 3.058 times higher in former users (p=0.029), 2.616 times (p=0.017) in low-users, and 3.547 times (p=0.001) higher in high-users.

Patients with complications were associated with an ASA score of 3 and above, 2.004 times (p=0.029) more than the ASA score I and II. In addition, the risk of developing complications is 2,323 times (p=0.027) higher in minimally invasive surgery and 4.458 times (p<0.001) in major surgery compared to minor surgery (Table 3).

Table 3: Evaluation of relationships of independent variables with development of complication using logistic regression analysis.

77 111	D.	Wald	P value	0.D*	CI† 95%	
Variables	В			OR*	Min	Max
Age groups (years)						
35-64 (ref. ≤35)	0.228	0.565	0.452	1.256	0.693	2.276
>65 (ref. ≤35)	0.519	1.671	0.196	1.680	0.765	3.688
Gender						
Male (ref.=female)	-0.067	0.032	0.857	0.935	0.451	1.941
Comorbidities		•		•	-	
+[ref.= (-)]	0.255	0.606	0.436	1.291	0.679	2.456
Tobacco use						
Former user (ref.=non-user)	1.118	4.753	0.029	3.058	1.120	8.353
Light user (ref.=non-user)	0.962	5.703	0.017	2.616	1.188	5.760
Heavy user (ref.=non-user)	1.266	11.998	0.001	3.547	1.733	7.260
ASA‡ score	0.695	4.770	0.029	2.004	1.074	3.740
Diagnosis						
Malign diseases (ref.=benign diseases)	-0.042	0.010	0.919	0.959	0.432	2.129
Infectious diseases (ref.=benign diseases)	0.084	0.034	0.853	1.088	0.445	2.657

Continued.

Variables	В	Wald	P value	OR*	CI† 95%	
variables	Б				Min	Max
Sp. pneumothorax (ref.=benign diseases)	0.683	3.690	0.055	1.980	0.986	3.974
Surgery types						
Minimally invasive (ref.=minor operations)	0.843	4.886	0.027	2.323	1.100	4.906
Major operations (ref.=minor operations)	1.495	31.670	< 0.001	4.458	2.649	7.502

Note: Hosmer-Lemeshow p=0.717, r²=0.177, *Odds Ratio, †Confidence interval, ‡American Society of Anesthesiologists.

Fifty (6.6%) of the patients were admitted to ITU for emergencies. Twenty-six (3.4%) of the patients had intubation, and 20 (2.6%) patients died. Fifty-five (7.3%) patients had at least one of these poor prognosis criteria. No significant relationship was found between tobacco use and ITU admission, intubation, and mortality. Logistic regression analysis was performed to reveal possible variables associated with poor prognosis factors (ITU admission, intubation, exitus).

Table 4 includes the relationship of independent variables in predicting the poor prognosis outcomes by logistic regression analysis. Poor prognostic outcomes were 5, 543 times (p=0.002) more common in those 65 years and older

than those under 35 years of age. Poor prognostic results were observed 2.015 times (p<0.001) more frequently in those with an ASA score of III and above than those with an ASA score below III (Table 4).

For all patients, the mean length of stay in hospital was 8.86 ± 6.95 days. In the short-stay group (8 days or less), 208 (48.3%) of 431 patients were using tobacco products, while 215 (66.6%) of 323 patients in the long-stay group (9 days or more) were using tobacco products. This result showed that the rate of tobacco use was significantly higher in the long-stay in hospital-group ($X^2=25.116$, p<0.001).

Table 4: Evaluation of the relationship of independent variables in the prediction of poor prognosis criteria (admission to intensive care unite-intubation-mortality) by logistic regression analysis.

Variables	В	Wald	P value	OR*	CI† 95%	
Variables					Min	Max
Age groups (years)						
35-64 (ref. ≤35)	0.836	2.483	0.115	2.308	0.815	6.534
>65 (ref. ≤35)	1.712	9.723	0.002	5.543	1.889	16.262
Gender						
Male (ref.=female)	-0.071	0.038	0.846	0.932	0.456	1.902
Comorbidities			•	•		
+[ref.= (-)]	-0.101	0.078	0.780	0.904	0.444	1.839
Tobacco use						
Former user (ref.=non-user)	-1.332	2.785	0.095	0.264	0.055	1.262
Light user (ref.=non-user)	0.082	0.030	0.864	1.086	0.425	2.774
Heavy user (ref.=non-user)	0.079	0.031	0.861	1.083	0.445	2.637
ASA‡ score	1.417	15.024	< 0.001	4.125	2.015	8.444
Diagnosis						
Malign diseases (ref.=benign diseases)	-0.076	0.040	0.842	0.927	0.439	1.955
Infectious diseases (ref.=benign diseases)	-0.773	1.371	0.242	0.462	0.127	1.684
Sp. pneumothorax (ref.=benign diseases)	-0.980	3.545	0.060	0.375	0.135	1.041
Surgery types						
Minimally invasive (ref.=minor operations)	-0.273	0.323	0.570	0.761	0.297	1.951
Major operations (ref.=minor operations)	-0.286	0.386	0.534	0.751	0.305	1.852

Note: Hosmer-Lemeshow p=0.528, r²=0.180, *Odds Ratio, †Confidence interval, ‡American Society of Anesthesiologists.

Table 5 includes the relationship between tobacco use and other possible independent variables with the length of stay in hospital by logistic regression analysis. The length of stay in hospital was 2.635 times higher in patients 65 years of age and over compared to patients under 35 years of age (p=0.001).

The length of stay in hospital was 1.948 times (p=0.016) higher in patients who used low amounts of tobacco

products than non-users and 3.706 times (p<0.001) higher in those who used high tobacco products than non-users. The length of stay in hospital was 2,839 times (p<0.001) higher in patients with an ASA score of III or more than those with an ASA score of less than III.

Infectious lung diseases increase the length of stay in hospital by 1.929 times (p=0.039) according to benign causes. In addition, the length of hospital stay was 5.063 times (p<0.001) higher in patients with minimally invasive

surgery than those who underwent minor surgery and 14.797 times (p<0.001) longer in patients who underwent

major surgery than those who underwent minor surgery (Table 5).

Table 5: Evaluation of the relationship of independent variables in predicting the length of stay in hospital with logistic regression analysis.

Vaniables	В	Wald	P value	OR*	CI† 95%	
Variables	ь				Min	Max
Age groups (years)						
35-64 (ref. ≤35)	0.383	2.628	0.105	1.466	0.923	2,329
>65 (ref. ≤35)	0.969	10.841	0.001	2.635	1.480	4,691
Gender						
Male (ref.=female)	0.139	0.333	0.564	1.149	0.717	1.842
Comorbidities		•	•			
+[ref.= (-)]	-0.011	0.002	0.964	0.989	0.605	1.615
Tobacco use						
Former user (ref.=non-user)	0.443	1.405	0.236	1.557	0.749	3.240
Light user (ref.=non-user)	0.667	5.763	0.016	1.948	1.130	3.358
Heavy user (ref.=non-user)	1.310	24.683	< 0.001	3.706	2.210	6.213
ASA‡ score	1.043	13.220	< 0.001	2.839	1.618	4.981
Diagnosis						
Malign diseases (ref.=benign diseases)	-0.073	0.070	0.792	0.929	0.539	1.602
Infectious diseases (ref.=benign diseases)	0.657	4.252	0.039	1.929	1.033	3.601
Sp. pneumothorax (ref.=benign diseases)	0.431	2.588	0.108	1.539	0.910	2.602
Surgery types						
Minimally invasive (ref.=minor operations)	1.622	32.826	< 0.001	5.063	2.907	8.817
Major operations (ref.=minor operations)	2.694	109.559	< 0.001	14.797	8.934	24.507

Note: Hosmer-Lemeshow p=0,372, r2=0,352, *Odds Ratio, †Confidence interval, ‡American Society of Anesthesiologists.

DISCUSSION

This study showed that tobacco use was associated with young age, male gender, development of postoperative complications, and high length of stay in hospital in thoracic surgery patients. However, according to our study, there was no relationship between tobacco use and ITU admission, intubation, and mortality.

According to Turkey's data of the Global Adult Tobacco Survey of the WHO (GATS), the tobacco use rate in Turkey was 31.6% in 2016.8 Among the cases hospitalized for pulmonary surgery in our study, the proportion of tobacco users was 56.1%. Accordingly, the rate of tobacco users was higher in the group hospitalized due to pulmonary surgical diseases.

According to Turkey's data, 44.1% of men in Turkey in 2016, 19.2% of women were using tobacco. In our study, 73.1% of men and 13.9% of women use tobacco. The rate of total smokers is higher than the general population. This difference is due to the higher tobacco use among men, as female tobacco use is similar to the general population. In our series, the higher rate of tobacco use in men compared to the general population may be attributed to the higher incidence of diseases such as primary spontaneous pneumothorax, which are more common in smoking men. 9

The mean age of smokers and non-smokers in different studies was at various rates. ¹⁰⁻¹² In these studies, smokers

are younger than non-smokers are. In our study, 62.8% of patients under the age of 65 were tobacco users. In one study, 63.9% of patients under the age of 65 were in the smoker group, and the findings are consistent with our results.¹³

Thoracic surgery disrupts respiratory functions in the postoperative period and increases the risk of postoperative complications.¹⁴ Thoracic surgery is an independent risk factor even, as it is performed directly to the respiratory system.¹⁵ The postoperative complication development rate after thorax surgery varies between 12% and 50%.¹⁶ In our study, complications developed in 12.73% of our patients. Most of the patients who developed complications were using tobacco products, and there was a statistically significant relationship between tobacco use and the rate of postoperative complications.

According to our study, cigarette increases the risk of postoperative pulmonary complications. However, there was no significant difference between cigarettes and other tobacco products related to the development of complications.

When previous studies are examined, different findings have been reported as postoperative pulmonary complications in each study. Findings frequently reported as postoperative complications in the literature; pneumonia and other respiratory infections, atelectasis, PAL, need for intubation, pneumothorax, respiratory

failure, pleural effusion, and bronchospasm. ¹⁷⁻¹⁹ The most common postoperative pulmonary complications in this study were PAL, atelectasis, pneumothorax, respiratory failure, pleural effusion and empyema, cardiac complications, surgical site infection, pneumonia, acute renal failure, and bronchospasm. These findings are complications in the literature at different rates.

Seok et al reported PAL and atelectasis as the most common postoperative pulmonary complications in their study. Similarly, the most common pulmonary complications in our study were PAL and atelectasis. PAL is defined as an air leak that lasts more than 5-7 days. There are studies evaluating smoking as a risk factor for PAL. Similarly, in our study, there was a significant relationship between PAL and tobacco use.

Atelectasis is a common complication in thoracic surgery clinics. It is more common in basal parts of the lung, and operated patients, especially in incisions close to the diaphragm.²³ Atelectasis developed in 3.18% of patients in our study, and it constituted 25% of patients who developed complications. There was a significant relationship between atelectasis and tobacco use.

When considered as a surgical complication, 2.65% of patients developed pneumothorax. One of the risk factors of spontaneous pneumothorax is tobacco use.²⁴ We found a statistically significant difference between tobacco use and the development of spontaneous pneumothorax as a complication. This result was consistent with the literature.

We did not find any significant relationship between other complications and tobacco use. However, since the number of these complications was low, we did not accept the results as reliable.

Logistic regression analysis was performed to evaluate possible factors associated with the development of complications together. Accordingly, the development of complications is significantly associated with tobacco use, ASA score, and type of surgery. The results of logistic regression analysis showed that pulmonary complications were seen more in former and current tobacco users compared to non-tobacco users. Major surgeries and minimally invasive surgeries are more associated with the development of complications, respectively than minor surgeries. The ASA score representing the physical performance of the patient was also found to be related to the development of complications.

The admission rate to ITU was significantly higher in patients over 65 years of age. This result is consistent with the results of the previous studies. However, contrary to some studies, no relationship was found between intensive therapy unit admission and smoking. The reason for this was that the patients using tobacco in our series were younger, whereas the patients who needed ITU were older.

We had to intubate 26 of the patients. The need for

intubation was not associated with tobacco use, there was a significant relationship between intubation need and age, and it revealed that older patients needed more intubation.

While some studies have shown that smoking increases mortality, some studies have shown no relationship between smoking and mortality. ²⁷⁻²⁹ Twenty patients died, 10 of whom used tobacco, and 10 of whom did not. Accordingly, there was no relationship between tobacco use and mortality. However, there was a significant relationship between age and death, and most patients who died were over 65 years old. This result was not surprising considering that tobacco users are mostly young. Ten of the patients who died had infectious diseases and empyema, seven had malignant diseases, and three had pleural effusion due to congestive heart failure. Eight patients had comorbidities.

There are close relationships between admission to the ITU, intubation, and death. Some patients were included in more than one of these groups. Therefore, these three groups were evaluated together in the logistic regression model as a poor prognosis group. According to this; the development of poor prognostic outcomes was significantly associated with age and ASA score (p <0.05). Poor physical performance and advanced age were associated with ITU admission, intubation, and death.

According to a study evaluating the relationship between hospital stay and smoking, the length of stay in hospital is higher in the smoker group.³⁰ In our study, the rate of tobacco use was higher in those who were hospitalized for more than 9 days. The relationship between the length of stay in hospital and tobacco use was consistent with the literature.

In order to evaluate the effect of smoking on the length of stay in hospital more accurately, the logistic regression analysis model was used together with other possible factors affecting the length of stay in hospital. According to logistic regression analysis results, Age, tobacco use, ASA score, diagnosis, and treatment methods were significantly related to the length of stay in hospital. According to the results of logistic regression analysis, the variable most associated with length of stay in hospital was the type of surgery. Major surgery and minimally invasive surgery, respectively, were more associated with length of stay in hospital than minor surgery. The patient group with a high length of stay in hospital had a higher proportion of current tobacco users. Heavy tobacco users had more length of stay in hospital than light tobacco users. Former tobacco users and non-tobacco users were not associated with the length of stay in hospital. The physical performance status of the patients was found to be related to the length of stay in hospital. The length of stay was higher in patients with ASA III and above. A poor physical condition prolonged hospital stay. Length of hospital stay was longer in infectious diseases. The reason for this was thought to be that antibiotic treatment was given in infectious diseases in the preoperative period and, if necessary, in the postoperative period. In addition, the length of stay in hospital was found to be higher in elderly patients compared to younger ages.

One limitation of this study was that it was a retrospective study, and therefore, the possibility of data loss. Another limitation is that only inpatients were included in the study. The strength of the study is that it was studied in three different big centers, and it was a large series.

CONCLUSION

In conclusion, the proportion of tobacco users was higher in patients with thoracic surgery than in those who did not. Tobacco use is associated with young age, postoperative complications, and length of hospital stay. Therefore, tobacco use is a severe health problem, especially at young ages. In the fight against tobacco, emphasizing that tobacco use will have surgical consequences and a high probability of complications after surgery and supporting tobacco cessation programs will contribute to public health and prevent economic loss.

ACKNOWLEDGEMENTS

This study was presented as an oral presentation in 9. International Trakya Family Medicine Congress and its abstract has been published in the congress abstract book.

Funding: No funding sources Conflict of interest: None declared

Ethical approval: The study was approved by the

Institutional Ethics Committee

REFERENCES

- 1. Ekerbicer HC, Berberoglu U, Inci MB. The struggle for a smoke-free environment: from past to today. Sakarya Tıp Dergisi. 2018;8(3):470-4.
- 2. Myles PS, Iacono GA, Hunt JO, Fletcher H, Morris J, McIlroy D, et al. Risk of respiratory complications and wound infection in patients undergoing ambulatory surgery: smokers versus nonsmokers. Anesthesiology. 2002;97(4):842-7.
- 3. Bilir N, Cakir B, Dagli E, Erguder T, Onder Z. Tobacco control policies in Turkey. WHO; Europe; 2009.
- 4. Kurucu S, Kartal M, Erenmemisoglu A. HPLJ analysis of Nicotiana rustica L. and chewing tobacco (Maras powder) alkaloids. FABAD J Pharm Sci.1998;23:63-4.
- Lechner AJ. Respiratory system terms and diseases. In: Lechner AJ, Matuschak GM, Brink DS, Gold S, eds. Respiratory: An integrated approach to disease. 1st ed. Turkey: Istanbul Medicine Bookstore; 2014: 3-8.
- 6. Moppett I, Curran J. Smoking and the surgical patient. British J Anaest. 2001;4(1):122-4.
- Mayhew D, Mendonca V, Murthy BVS. A review of ASA physical status- historical perspectives and

- modern developments. Anaesthesia. 2019;74(3):373-9.
- 8. WHO. Global Adult Tobacco Survey (GATS), fact sheet Turkey, 2016-17. Available at: https://www.who.int/tobacco/surveillance/survey/gats/GATS_India_2016-17_FactSheet. Accessed on 10 May 2021.
- 9. Noppen M. Spontaneous pneumothorax: epidemiology, pathophysiology and cause. Eur Respir Rev. 2010;117(19):217-9.
- 10. Musallam K, Rosendaal FR, Zaatari G, Soweid A, Hoballah JJ, Sfeir PM, et al. Smoking and risk of mortality and vascular and respiratory events in patients undergoing major surgery. JAMA Surg. 2013;148(8):755-62.
- 11. Hawn MT, Houston TK, Campagna EJ, Graham LA, Singh J, Bishop M, et al. The attributable risk of smoking on surgical complications. Ann Surg. 2011;254(6):914-20.
- 12. Lugg ST, Tikka T, Agostini PJ, Kerr A, Adams K, Kalkat MS, et al. Smoking and timing of cessation on postoperative pulmonary complications after curative-intent lung cancer surgery. J Cardiothorac Surg. 2017;12(1):52.
- 13. Moller AM, Maaløe R, Pedersen T. Postoperative intensive care admittance: The role of tobacco smoking. Acta Anaesthesiol Scand. 2001;45(3):345-8.
- 14. Agostini P, Cieslik H, Rathinam S, Bishay A, Kalkat MS, Rajesh PB, et al. Postoperative pulmonary complications following thoracic surgery: are there any modifiable risk factors? Thorax. 2010;65(9):815-8.
- 15. Nakagawa M, Tanaka H, Tsukuma H, Kishi Y. The preoperative smoke-free period and the incidence of postoperative pulmonary complications after pulmonary surgery. Chest. 2001;120(3):705-10.
- 16. Saad IAB, Capitani EM, Toro IFC, Zambon L. Clinical variables of preoperative risk in thoracic surgery. Sao Paulo Med J. 2003;121(3):107-10.
- 17. Seok Y, Hong N, Lee E. Impact of smoking history on postoperative pulmonary complications: a review of recent lung cancer patients. Ann Thorac Cardiovasc Surg. 2014;20(2):123-8.
- 18. Turan A, Mascha EJ, Roberman D, Turner PL, You J, Kurz A, et al. Smoking and perioperative outcomes. Anesthesiology. 2011;114(4):837-46.
- 19. Jammer I, Wickboldt N, Sander M, Smith A, Schults MJ, Pelosi P, et al. Standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions. Eur J Anaesthesiol. 2015;32(2):88-105.
- 20. Miskovic A, Lumb AB. Postoperative pulmonary complications. Br J Anaesth. 2017 1;118(3):317-34.
- 21. Dugan KC, Laxmanan B, Murgu S, Hogarth K. Management of persistent air leaks. Chest 2017;152(2):417-23.
- 22. Liberman M, Muzikansky A, Wright CD, Wain JC, Donehue DM, Allan JS, et al. Incidence and risk

- factors of persistent air leak after major pulmonary resection and use of chemical pleurodesis. Ann Thorac Surg. 2010;89(3):891-8.
- 23. Rudra A, Das S. Postoperative pulmonary complications. Indian J Anaesth. 2006;50(2):89-98.
- 24. Bintcliffe O, Maskell N. Spontaneous pneumothorax. BMJ. 2014;348:2928.
- 25. Sakata KK, Reisenauer JS, Kern RM, Mullon JJ. Persistent air leak-review. Respir Med. 2018;137:213-8.
- Gronkjær M, Eliasen M, Ettrup LS, Tolstrup JS, Christiansen AH, Mikkelsen SS, et al. Preoperative smoking status and postoperative complications: a systematic review and meta-analysis. Ann Surg. 2014;259(1):52-71.
- 27. Myers K, Hajek P, Hinds C, McRobbie H. Stopping smoking shortly before surgery and postoperative complications. Arch Intern Med. 2011;171(11):983-9.

- 28. Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO. Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. Am J Med. 2011;124(2):144-54.
- Sarraf N, Thalib L, Hughes A, Tolan M, Young V, McGovern E. Effect of smoking on short-term outcome of patients undergoing coronary artery bypass surgery. Ann Thorac Surg. 2008;86(2):517-23.
- 30. Rezaei S, Akbari Sari A, Arab M, Majdzadeh R, Shaahmadi F, et al. The association between smoking status and hospital length of stay: evidence from a hospital-based cohort. Hosp Pract. 2016;44(3):129-32.

Cite this article as: Degirmenci M, Kus C. Evaluation of post-operative complications associated with tobacco use in thoracic surgery patients. Int Surg J 2021;8:2272-80.